

## US 7,300,662 B2

Page 3

5,376,112 A	12/1994	Duran	5,591,224 A	1/1997	Schwartz et al.
5,378,475 A	1/1995	Smith et al. .... 424/473	5,591,227 A	1/1997	Dinh et al.
5,380,299 A	1/1995	Fearnott et al.	5,599,352 A	2/1997	Dinh et al.
5,382,261 A	1/1995	Palmar	5,603,722 A	2/1997	Phan et al.
5,383,853 A	1/1995	Jung et al. .... 604/103.04	5,604,283 A	2/1997	Wada et al. .... 524/236
5,383,928 A	1/1995	Scott et al.	5,605,696 A	2/1997	Eury et al.
5,387,235 A	2/1995	Chuter	5,607,463 A	3/1997	Schwartz et al.
5,389,106 A	2/1995	Tower	5,607,475 A	3/1997	Cahalan et al.
5,391,730 A	2/1995	Skotnicki et al. .... 540/456	5,609,629 A	3/1997	Fearnott et al.
5,393,772 A	2/1995	Yue et al.	5,616,608 A	4/1997	Kinsella et al. .... 514/449
5,395,390 A	3/1995	Simon et al.	5,620,984 A	4/1997	Bianco et al.
5,397,355 A	3/1995	Marin et al.	5,621,102 A	4/1997	Bianco et al.
5,399,352 A	3/1995	Hanson ..... 424/423	5,622,975 A	4/1997	Singh et al.
5,403,341 A	4/1995	Solar	5,624,411 A	4/1997	Tuch
5,405,377 A	4/1995	Cragg	5,628,785 A	5/1997	Schwartz et al.
5,409,696 A	4/1995	Narayanan et al.	5,629,077 A	5/1997	Turnlund et al.
5,411,549 A	5/1995	Peters	5,629,315 A	5/1997	Bianco et al.
5,415,619 A	5/1995	Lee et al.	5,632,763 A	5/1997	Glastra
5,417,969 A	5/1995	Hsu et al. .... 424/78.27	5,632,771 A	5/1997	Boatman et al. .... 623/1.15
5,419,760 A	5/1995	Narciso, Jr.	5,632,776 A	5/1997	Kurumatani et al. .... 424/423
D359,802 S	6/1995	Fontaine	5,632,840 A	5/1997	Campbell
5,421,955 A	6/1995	Lau	5,635,201 A	6/1997	Fabo ..... 424/443
5,423,885 A	6/1995	Williams	5,637,113 A	6/1997	Tartaglia et al.
5,429,618 A	7/1995	Keogh	5,643,312 A	7/1997	Fischell et al.
5,429,634 A	7/1995	Narciso	5,643,939 A	7/1997	Ohlstein
5,439,446 A	8/1995	Barry	5,646,160 A	7/1997	Morris et al.
5,441,515 A	8/1995	Khosravi et al.	5,648,357 A	7/1997	Bianco et al.
5,441,516 A	8/1995	Wang et al.	5,649,952 A	7/1997	Lam
5,441,947 A	8/1995	Dodge et al.	5,649,977 A	7/1997	Campbell
5,443,458 A	8/1995	Eury	5,651,174 A	7/1997	Schwartz et al.
5,443,477 A	8/1995	Marin et al.	5,652,243 A	7/1997	Bianco et al.
5,443,496 A	8/1995	Schwartz et al.	5,653,747 A	8/1997	Dereume ..... 623/1.54
5,443,498 A	8/1995	Fontaine	5,653,992 A	8/1997	Bezswana et al.
5,443,500 A	8/1995	Sigwart	5,662,609 A	9/1997	Slepian
5,447,724 A	9/1995	Heimus et al.	5,665,591 A	9/1997	Sonenshein et al. .... 435/375
5,449,372 A	9/1995	Schmaltz et al.	5,665,728 A	9/1997	Morris et al.
5,449,373 A	9/1995	Pinchasik et al.	5,667,764 A	9/1997	Kopia et al. .... 424/1.45
5,449,382 A	9/1995	Dayton	5,669,924 A	9/1997	Shaknovich
5,464,450 A	11/1995	Buscemi et al.	5,670,506 A	9/1997	Leigh et al.
5,464,540 A	11/1995	Friesen et al. .... 210/640	5,672,638 A	9/1997	Verhoeven et al. .... 523/112
5,464,650 A	11/1995	Berg et al.	5,674,242 A	10/1997	Phan et al. .... 606/198
5,474,563 A	12/1995	Myler et al. .... 606/108	5,679,400 A	10/1997	Tuch
5,486,357 A	1/1996	Narayanan	5,679,659 A	10/1997	Verhoeven et al.
5,496,365 A	3/1996	Sgro	5,684,061 A	11/1997	Ohnishi et al. .... 523/114
5,500,013 A	3/1996	Buscemi et al.	5,691,311 A	11/1997	Maraganore et al. .... 514/12
5,504,091 A	4/1996	Molnar-Kimber et al. ... 514/291	5,693,085 A	12/1997	Buirge et al.
5,510,077 A	4/1996	Dinh et al.	5,697,967 A	12/1997	Dinh et al.
5,512,055 A	4/1996	Domb et al. .... 604/265	5,697,971 A	12/1997	Fischell et al.
5,516,781 A	5/1996	Morris et al.	5,700,286 A	12/1997	Tartaglia et al.
5,519,042 A	5/1996	Morris et al. .... 514/378	5,707,385 A	1/1998	Williams
5,523,092 A	6/1996	Hanson et al.	5,709,874 A	1/1998	Hanson et al.
5,527,354 A	6/1996	Fontaine et al.	5,713,949 A	2/1998	Jayaraman ..... 623/1.12
5,545,208 A	8/1996	Wolff et al.	5,716,981 A	2/1998	Hunter et al. .... 514/449
5,551,954 A	9/1996	Buscemi et al.	5,725,549 A	3/1998	Lam
5,554,182 A	9/1996	Dinh et al.	5,725,567 A	3/1998	Wolff et al.
5,554,954 A	9/1996	Takahashi	5,728,150 A	3/1998	McDonald et al.
5,556,413 A	9/1996	Lam	5,728,420 A	3/1998	Keogh
5,559,122 A	9/1996	Nelson et al. .... 514/291	5,731,326 A	3/1998	Hart et al.
5,562,922 A	10/1996	Lambert	5,733,327 A	3/1998	Igaki et al.
5,563,146 A	10/1996	Morris et al.	5,733,920 A	3/1998	Mansuri et al.
5,569,197 A	10/1996	Helmus et al.	5,733,925 A	3/1998	Kunz et al.
5,569,295 A	10/1996	Lam	5,735,897 A	4/1998	Buirge
5,569,462 A	10/1996	Martinson et al. .... 424/423	5,739,138 A	4/1998	Bianco et al.
5,569,463 A	10/1996	Helmus et al. .... 424/426	5,755,734 A	5/1998	Richter et al.
5,571,089 A	11/1996	Crocker ..... 604/103.01	5,755,772 A	5/1998	Evans et al. .... 128/898
5,571,166 A	11/1996	Dinh et al.	5,759,205 A	6/1998	Valentini ..... 433/173
5,574,059 A	11/1996	Regunathan et al.	5,769,883 A	6/1998	Buscemi et al.
5,575,818 A	11/1996	Pinchuk ..... 623/1.15	5,776,184 A	7/1998	Tuch
5,578,075 A	11/1996	Dayton	5,780,462 A	7/1998	Lee et al. .... 514/183
5,580,873 A	12/1996	Bianco et al.	5,780,476 A	7/1998	Underiner et al.
5,580,874 A	12/1996	Bianco et al.	5,782,908 A	7/1998	Cahalan et al.
5,591,140 A	1/1997	Narayanan et al.	5,788,979 A	8/1998	Alt
5,591,197 A	1/1997	Orth et al.	5,792,106 A	8/1998	Mische ..... 604/103.01

## US 7,300,662 B2

Page 4

5,792,772 A	8/1998	Bianco et al.	6,179,817 B1	1/2001	Zhong	604/265
5,798,372 A	8/1998	Davies et al.	6,187,757 B1	2/2001	Clackson et al.	514/31
5,799,384 A	9/1998	Schwartz et al.	6,193,746 B1	2/2001	Strecker	623/1.13
5,800,507 A	9/1998	Schwartz	6,214,901 B1	4/2001	Chudzik et al.	
5,800,508 A	9/1998	Goicoechea et al.	6,225,346 B1	5/2001	Tang et al.	514/523
5,807,861 A	9/1998	Klein et al.	6,240,616 B1	6/2001	Yan	
5,811,447 A	9/1998	Kunz et al.	6,245,537 B1	6/2001	Williams et al.	435/135
5,820,917 A	10/1998	Tuch	6,251,920 B1	6/2001	Grainger et al.	514/319
5,820,918 A	10/1998	Ronan et al.	6,254,632 B1	7/2001	Wu et al.	
5,824,048 A	10/1998	Tuch	6,254,634 B1	7/2001	Anderson et al.	623/1.42
5,824,049 A	10/1998	Ragheb et al.	6,258,121 B1	7/2001	Yang et al.	
5,827,587 A	10/1998	Fukushi	6,268,390 B1	7/2001	Kunz	
5,833,651 A	11/1998	Donovan et al.	6,273,913 B1	8/2001	Wright et al.	
5,837,008 A	11/1998	Berg et al.	6,284,305 B1	9/2001	Ding et al.	427/2.28
5,837,313 A	11/1998	Ding et al.	6,287,320 B1	9/2001	Slepian	
5,843,120 A	12/1998	Israel et al.	6,287,628 B1	9/2001	Hossainy et al.	
5,843,166 A	12/1998	Lentz et al.	6,299,604 B1	10/2001	Ragheb et al.	
5,843,172 A	12/1998	Yan	6,306,144 B1	10/2001	Sydney et al.	606/108
5,849,034 A	12/1998	Schwartz	6,306,166 B1	10/2001	Barry et al.	
5,851,217 A	12/1998	Wolff et al.	6,306,176 B1	10/2001	Whitbourne	
5,851,231 A	12/1998	Wolff et al.	6,306,421 B1	10/2001	Kunz et al.	
5,858,990 A	1/1999	Walsh	6,309,380 B1	10/2001	Larson et al.	
5,861,027 A	1/1999	Trapp	6,309,660 B1	10/2001	Hsu et al.	
5,865,814 A	2/1999	Tuch	6,313,264 B1	11/2001	Caggiano et al.	
5,871,535 A	2/1999	Wolff et al.	6,316,018 B1	11/2001	Ding et al.	424/423
5,873,904 A	2/1999	Ragheb et al.	6,335,029 B1	1/2002	Kamath et al.	424/423
5,876,433 A	3/1999	Lunn	6,358,556 B1	3/2002	Ding et al.	427/2.24
5,877,224 A	3/1999	Brocchini et al.	6,369,039 B1	4/2002	Palasis et al.	424/93.2
5,879,697 A	3/1999	Ding et al.	6,379,382 B1	4/2002	Yang	623/1.42
5,882,335 A	3/1999	Leone et al.	6,387,121 B1	5/2002	Alt	623/1.15
5,891,108 A	4/1999	Leone et al.	6,403,635 B1	6/2002	Kinsella et al.	514/449
5,893,840 A	4/1999	Hull et al.	6,407,067 B1	6/2002	Schafer	514/19
5,897,911 A	4/1999	Loeffler	6,517,858 B1	2/2003	Le Moel et al.	424/424
5,900,246 A	5/1999	Lambert	6,517,889 B1	2/2003	Jayaraman	427/2.24
5,902,266 A	5/1999	Leone et al.	6,545,097 B2	4/2003	Pinchuk et al.	525/240
5,912,253 A	6/1999	Cottens et al.	6,585,764 B2	7/2003	Wright et al.	623/1.42
5,916,910 A	6/1999	Lai	6,620,194 B2	9/2003	Ding et al.	623/1.43
5,922,393 A	7/1999	Jayaraman	6,746,773 B2	6/2004	Llanos et al.	428/421
5,922,730 A	7/1999	Hu et al.	6,776,796 B2	8/2004	Falotico et al.	623/1.46
5,932,243 A	8/1999	Fricker et al.	6,808,536 B2	10/2004	Wright et al.	623/1.42
5,932,299 A	8/1999	Katoot	2001/0007083 A1	7/2001	Roorda	
5,932,580 A	8/1999	Levitzi et al.	2001/0029351 A1	10/2001	Falotico et al.	604/103.02
5,951,586 A	9/1999	Berg et al.	2001/0029660 A1	10/2001	Johnson	
5,957,971 A	9/1999	Schwartz	2001/0032014 A1	10/2001	Yang et al.	
5,968,091 A	10/1999	Pinchuk et al.	2001/0034363 A1	10/2001	Li et al.	
5,972,027 A	10/1999	Johnson	2001/0037145 A1	11/2001	Guruwaiya et al.	
5,976,534 A	11/1999	Hart et al.	2002/0010418 A1	1/2002	Lary et al.	604/101.04
5,977,163 A	11/1999	Li et al.	2002/0032477 A1	3/2002	Helmus et al.	623/1.2
5,980,553 A	11/1999	Gray et al.	2002/0041899 A1	4/2002	Chudzik et al.	424/487
5,980,566 A	11/1999	Alt et al.	2002/0061326 A1	5/2002	Li et al.	424/424
5,980,972 A	11/1999	Ding	2002/0068969 A1	6/2002	Shanley et al.	623/1.16
5,981,568 A	11/1999	Kunz et al.	2002/0071902 A1	6/2002	Ding et al.	427/2.24
5,985,307 A	11/1999	Hanson et al.	2002/0082680 A1	6/2002	Shanley et al.	623/1.16
5,997,468 A	12/1999	Wolff et al.	2002/0082685 A1	6/2002	Sirhan et al.	623/1.42
6,004,346 A	12/1999	Wolff et al.	2002/0091433 A1	7/2002	Ding et al.	623/1.2
6,015,432 A	1/2000	Rakos et al.	2002/0095114 A1	7/2002	Palasis	604/96.01
6,015,815 A	1/2000	Mollison	2002/0099438 A1	7/2002	Furst	623/1.16
6,039,721 A	3/2000	Johnson et al.	2002/0103526 A1	8/2002	Steinke	623/1.11
6,059,813 A	5/2000	Vrba et al.	2002/0119178 A1	8/2002	Levesque et al.	424/423
6,071,305 A	6/2000	Brown et al.	2002/0123505 A1	9/2002	Mollison et al.	514/291
6,074,659 A	6/2000	Kunz et al.	2002/0127327 A1	9/2002	Schwartz et al.	427/2.15
6,080,190 A	6/2000	Schwartz	2002/0133222 A1	9/2002	Das	623/1.16
6,096,070 A	8/2000	Ragheb et al.	2002/0133224 A1	9/2002	Bajgar et al.	623/1.39
6,120,536 A	9/2000	Dinge et al.	2002/0165608 A1	11/2002	Llanos	604/500
6,120,847 A	9/2000	Yang et al.	2002/0193475 A1	12/2002	Hossainy et al.	524/113
6,136,798 A	10/2000	Cody et al.	2003/0065377 A1	4/2003	Davila et al.	604/265
6,140,127 A	10/2000	Sprague	2003/0216699 A1	11/2003	Falotico	
6,146,358 A	11/2000	Rowe	2004/0049265 A1	3/2004	Ding et al.	623/1.42
6,153,252 A	11/2000	Hossainy et al.	2004/0243097 A1	12/2004	Falotico et al.	604/500
6,159,488 A	12/2000	Nagier et al.	2004/0260268 A1	12/2004	Falotico et al.	604/500
6,171,232 B1	1/2001	Papandreu et al.	2005/0002986 A1	1/2005	Falotico et al.	424/426
6,171,609 B1	1/2001	Kunz	2005/0004663 A1	1/2005	Llanos et al.	623/1.46
6,177,272 B1	1/2001	Nabel et al.	2005/0033261 A1	2/2005	Falotico et al.	604/500

A1237

## US 7,300,662 B2

Page 5

2005/0106210 A1	5/2005	Ding et al.	424/423
2005/0187611 A1	8/2005	Ding et al.	623/1.15
2005/0208200 A1	9/2005	Ding et al.	427/2.25
2006/0088654 A1	4/2006	Ding et al.	427/2.21
2006/0089705 A1	4/2006	Ding et al.	623/1.15

## FOREIGN PATENT DOCUMENTS

DE	19723723 A1	12/1998
EP	0 145 166 A2	6/1985
EP	0 177 330 A2	4/1986
EP	0 183 372 A1	6/1986
EP	0 221 570 A2	5/1987
EP	0 421 729 A2	4/1991
EP	0 540 290 A2	10/1992
EP	0 568 310 A1	11/1993
EP	0 604 022 A1	6/1994
EP	0 621 015 A1	10/1994
EP	0 623 354 A1	11/1994
EP	0 734 698 A2	3/1996
EP	0 712 615 A1	5/1996
EP	0 716 836 A1	6/1996
EP	0 800 801 A1	8/1996
EP	0 734 721 A1	10/1996
EP	0 747 069 A2	12/1996
EP	0 761 251 A1	3/1997
EP	0 830 853 A1	7/1997
EP	0 540 290 B1	1/1998
EP	0 815 803 A1	7/1998
EP	0 850 651 A2	7/1998
EP	0 938 878 A3	9/1999
EP	0950386 B1	10/1999
EP	0 968 688 A1	1/2000
EP	0 633 032 B1	2/2001
EP	1 192 957 A2	4/2002
EP	1 588 726 A1	10/2005
EP	1 588 727 A1	10/2005
FR	0 566 807 A1	4/1992
GB	1 205 743	9/1970
GB	2 135 585 A	9/1984
GB	0 662 307 A2	12/1994
SU	660689	5/1979
SU	1457921	2/1989
WO	89/03232 A1	4/1989
WO	WO 91/12779 A1	9/1991
WO	WO 92/15286 A1	9/1992
WO	WO 94/01056 A1	1/1994
WO	WO 94/21308 A1	9/1994
WO	WO 94/21309 A1	9/1994
WO	WO 94/24961 A1	11/1994
WO	WO 96/00272 A1	1/1996
WO	WO 96/26689 A1	9/1996
WO	WO 96/32907 A1	10/1996
WO	WO 96/34580 A1	11/1996
WO	96/41807 A1	12/1996
WO	WO 97/25000 A1	7/1997
WO	WO 97/33534 A1	9/1997
WO	97/35575 A1	10/1997
WO	98/08463 A1	3/1998
WO	WO 98/13344 A1	4/1998
WO	WO 98/19628 A1	5/1998
WO	98/23244 A1	6/1998
WO	WO 98/23228 A1	6/1998
WO	WO 98/34669 A1	8/1998
WO	WO 98/36784 A1	8/1998
WO	WO 98/47447 A1	10/1998
WO	WO 98/56312 A1	12/1998
WO	00/21584 A1	4/2000
WO	00/27455 A1	5/2000
WO	WO 00/27445 A1	5/2000
WO	00/32255 A1	6/2000
WO	00/38754 A1	7/2000
WO	01/87342 A2	11/2001

WO	01/87372 A1	11/2001
WO	01/87373 A1	11/2001
WO	WO 01/87376 A1	11/2001
WO	02/26139 A1	4/2002
WO	02/26271 A1	4/2002
WO	02/26280 A1	4/2002
WO	02/26281 A1	4/2002
WO	03/015664 A1	2/2003
WO	03/057218 A1	7/2003

## OTHER PUBLICATIONS

U.S. Appl. No. 07/819,314, filed Jan. 9, 1992, Morris.

U.S. Appl. No. 08/424,884, filed Apr. 19, 1995, Helmus et al.

U.S. Appl. No. 08/526,273, filed Sep. 11, 1995, Ding.

U.S. Appl. No. 08/730,542, filed Oct. 11, 1996, Helmus.

U.S. Appl. No. 09/575,480, filed May 19, 2000, Kopia.

U.S. Appl. No. 10/431,059, filed May 7, 2003, Falotico.

Abraham, R. T., "Mammalian target of rapamycin: Immunosuppressive drugs offer new insight into cell growth regulation," *Progress in Inflammation Research*, 2000, Switzerland.

Alvarado, R. et al., "Evaluation of Polymer-coated Balloon-expandable Stents in Bile Ducts," *Radiology*, 1989, 170, 975-978.

Badimon, J. J. et al., "Inhibitory Effects of Rapamycin on Intimal Hyperplasia After PTCA," *JACC*, Mar. 1998.

Bailey et al., "Polymer Coating of Palmaz-Schatz Stent Attenuates Vascular Spasm after Stent Placement," *Circulation*, 82:III-541 (1990).

Berk, B. C. et al., "Pharmacologic Roles of Heparin and Glucocorticoids to Prevent Restenosis After Coronary Angioplasty," *JACC*, May 1991, 17(6), 111B-117B.

Bertram, P. G. et al., "The 14-3-3 proteins positively regulate rapamycin-sensitive signaling," *Current Biology*, 1998, 8, 1259-1267.

Biomaterials Science (B.D. Ratner, Ed.), Academic Press, New York, NY, pp. 228-238, 1996.

Campbell, G. R. et al., "Phenotypic Modulation of Smooth Muscle Cells in Primary Culture, Vascular Smooth Muscle Cells in Culture," *CRC Press*, 1987, 39-55.

Chang, M. W. et al., "Adenovirus-mediated Over-expression of the Cyclin/Cyclin-dependent Kinase inhibitor, p21 inhibits Vascular Smooth Muscle Cell Proliferation and Neointima Formation in the Rat Carotid Artery Model of Balloon Angioplasty," *J. Clin. Invest.*, 1995, 96, 2260-2268.

Chung, J. et al., "Rapamycin-FKBP specifically blocks growth-dependent activation of and signaling by the 70 kd S6 protein kinases," *Cell*, Jun. 26, 1992, 69(7), 1227-1236.

Clowes, A. W. et al., "Kinetics of cellular proliferation after arterial injury. IV. Heparin inhibits rat smooth muscle mitogenesis and migration," *Circ. Res.*, 1986, 58(6), 839-845.

Clowes, A. W. et al., Kinetics of Cellular Proliferation after Arterial Injury, *Laboratory Investigation*, 1985, 52(6), 611-616.

Clowes, A. W. et al., "Significance of quiescent smooth muscle migration in the injured rat carotid artery," *Circ Res.* 1985, 56(1), 139-145.

Clowes, A. W., "Suppression by heparin of smooth muscle cell proliferation in injured arteries," *Nature*, 1977, 265(5595), 625-626.

Colburn, M. D. et al., "Dose responsive suppression of myointimal hyperplasia by dexamethasone," *J. Vasc. Surg.*, 1992, 15, 510-518.

Currier, J. W. et al., "Colchicine Inhibits Restenosis After Iliac Angioplasty in the Atherosclerotic Rabbit," *Circ.*, 1989, 80(4), 11-66 (Abstract No. 0263).

Encyclopedia of Polymer Science and Engineering, vol. 7, Fluococarbon Elastomers, p. 257-267, Mar. 1989.

Farb, A. et al., "Vascular smooth muscle cell cytotoxicity and sustained inhibition of neointimal formation by fibroblast growth factor 2-saporin fusion protein," *Circ. Res.*, 1997, 80, 542-550.

Ferns, G. A. A. et al., "Inhibition of Neointimal Smooth Muscle Accumulation After Angioplasty by an Antibody to PDGF," *Science*, 1991, 253, 1129-1132.

US 7,300,662 B2

Page 6

- Fischman, D. L. et al., "A Randomized Comparison of Coronary Stent Placement and Balloon Angioplasty in the Treatment of Coronary Artery Disease," *N. Eng. J. Med.*, Aug. 25, 1994, 331(8), 496-501.
- Franklin, S. M. et al., "Pharmacologic prevention of restenosis after coronary angioplasty: review of the randomized clinical trials," *Coronary Artery Disease* Mar. 1993, 4(3), 232-242.
- Fukuyama, J. et al., "Tranilast suppresses the vascular intimal hyperplasia after balloon injury in rabbits fed on a high-cholesterol diet," *Eur. J. Pharmacol.*, 1996, 318, 327-332.
- Gregory, C. R. et al., "Rapamycin Inhibits Arterial Intimal Thickening Caused by Both Alloimmune and Mechanical Injury," *Transplantation*, Jun. 1993, 55(6), 1409-1418.
- Gregory, C. R. et al., "Treatment with Rapamycin and Mycophenolic Acid Reduces Arterial Intimal Thickening Produced by Mechanical Injury and Allows Endothelial Replacement," *Transplantation*, Mar. 15, 1995, 59(5), 655-661.
- Guyton, J. R. et al., "Inhibition of rat arterial smooth muscle cell proliferation by heparin. In vivo studies with anticoagulant and nonanticoagulant heparin," *Circ. Res.*, 1980, 46, 625-634.
- Hansson, G. K. et al., "Interferon- $\gamma$  Inhibits Arterial Stenosis After Injury," *Circ.*, 1991, 84, 1266-1272.
- Hashemolhosseini, S. et al., "Rapamycin Inhibition of the G1 to S Transition Is Mediated by Effects on Cyclin D1 mRNA and Protein Stability," *J. Biol. Chem.*, Jun. 5, 1998, 273, 14424-14429.
- Jonasson, J. et al., "Cyclosporin A inhibits smooth muscle proliferation in the vascular response to injury," *Proc. Natl. Acad. Sci.*, 1988, 85, 2303-2306.
- Kuhnt, M. et al., "Microbial Conversion of Rapamycin," *Enzyme and Microbial Technology*, 1997, 21, 405-412.
- Lange, R. A. MD et al., "Restenosis After Coronary Balloon Angioplasty," *Annu. Rev. Med.*, 1991, 42, 127-132.
- Liu, M. W. et al., "Trapidil in Preventing Restenosis After Balloon Angioplasty in the Atherosclerotic Rabbit," *Circ.*, 1990, 81, 1089-1093.
- Liu, M. W., MD et al., "Restenosis After Coronary Angioplasty Potential Biologic Determinants and Role of Intimal Hyperplasia," *Circulation*, 1989, 79, 1374-1387.
- Lundergan, C. F. et al., "Peptide inhibition of Myointimal Proliferation by Angiopeptin, a Somatostatin Analogue," *JACC*, May 1991, 17(6), 132B-136B.
- Majesky, M. W. et al., "Heparin regulates smooth muscle S phase entry in the injured rat carotid artery," *Circ. Res.*, 1987, 61, 296-300.
- Marx, S. O. et al., "Rapamycin-FKBP Inhibits Cell Cycle Regulators of Proliferation in Vascular Smooth Muscle Cells," *Circ. Res.*, 1995, 76, 412-417.
- Nemecsek, G. M. et al., "Terbinafine Inhibits the Mitogenic Response to Platelet-Derived Growth Factor in Vitro and Neoinimal Proliferation in Vivo," *J. Pharmacol. Exp. Ther.*, 1989, 248, 1167-1174.
- Okada, T. et al., "Localized Release of Perivascular Heparin Inhibits Intimal Proliferation after Endothelial Injury without Systemic Anticoagulation," *Neurosurgery*, 1989, 25, 892-898.
- Poon, M. et al., "Rapamycin Inhibits Vascular Smooth Muscle Cell Migration," *J. Clin. Invest.*, Nov. 1996, 98(10), 2277-2283.
- Popma, J. J. et al., "Clinical trials of restenosis after coronary angioplasty," *Circulation*, Sep. 1991, 84(3), 1426-1436.
- Powell, J. S. et al., "Inhibitors of Angiotensin-Converting Enzyme Prevent Myointimal Proliferation After Vascular Injury," *Science*, 1989, 245, 186-188.
- Rensing, B. J. et al., "Coronary restenosis elimination with a sirolimus eluting stent," *European Heart Journal*, 2001, 22, 2125-2130.
- Rodeck, C. et al., "Methods for the Transcervical Collection of Fetal Cells During the First Trimester of Pregnancy," *Prenatal Diagnosis*, 1995, 15, 933-942.
- Serruys, P. W. et al., "A comparison of balloon-expandable-stent implantation with balloon angioplasty in patients with coronary artery disease," *N Engl J Med*, Aug. 25, 1994; 331(8), 489-495.
- Serruys, P. W. et al., "Evaluation of ketanserine in the prevention of restenosis after percutaneous transluminal coronary angioplasty. A multicenter randomized double-blind placebo-controlled trial," *Circulation*. Oct. 1993; 88(4 Pt 1), 1588-1601.
- Serruys, P. W. et al., "Heparin-coated Palmaz-Schatz stents in human coronary arteries. Early outcome of the Benestent-II Pilot Study," *Circulation*, Feb. 1, 1996; 93(3), 412-422.
- Siekierka, J. J., "Probing T-Cell Signal Transduction Pathways with the Immunosuppressive Drugs, FK-506 and Rapamycin," *Immunologic Research*, 1994, 13, 110-116.
- Sigwart, et al., "Intravascular Stents to Prevent Occlusion and Restenosis After Transluminal Angioplasty," *N. Engl. J. Med.*, Mar. 19, 1987, 316, 701-706.
- Simons, M. et al., "Antisense c-myc oligonucleotides inhibit intimal arterial smooth muscle cell accumulation in vivo," *Nature*, 1992, 359, 67-70.
- Snow, A. D. et al., "Heparin modulates the composition of the extracellular matrix domain surrounding arterial smooth muscle cells," *Am. J. Pathol.*, 1990, 137, 313-330.
- Sollott, S. J. et al., "Taxol Inhibits Neointimal Smooth Muscle Cell Accumulation after Angioplasty in the Rat," *J. Clin. Invest.*, 1995, 95, 1869-1876.
- van Der Giessen, et al., "Self-expandable Mesh Stents: an Experimental Study Comparing Polymer Coated and Uncoated Wallstent Stents in the Coronary Circulation of Pigs," *Circulation* 1990, 82(suppl. III):III-542.
- van Der Giessen, W. J. et al., "Coronary stenting with polymer-coated and uncoated self-expanding endoprostheses in pigs," *Coron. Art. Disease* 1992; 3, 631-640.
- Vasey, C. G. et al., "Clinical Cardiology: Stress Echo and Coronary Flow", *Circulation*, Oct. 1989, 80(4) Supplement II, II-66.
- Verweire, E. et al., "Evaluation of Fluorinated Polymers As Coronary Stent Coating," *Journal of Materials Science: Materials in medicine*, Apr. 2000.
- Weinberger, J. et al., "Intracoronary irradiation: dose response for the prevention of restenosis in swine," *Int. J. Rad. Onc. Biol. Phys.*, 1996, 36, 767-775.
- Preliminary Amendment in U.S. Appl. No. 07/258,189, May 22, 1989.
- Trial Transcript from Nov. 6, 2000 at 185-90 and 235-36 (Attorney's opening remarks regarding '984 patent).
- Trial Transcript from Nov. 7, 2000 at 274-301, 307-315, 320-28 and 332 (Cordis expert testimony regarding the Palmaz-Schatz stent); 370-379, 480-496 (J. Palmaz testimony regarding the Palmaz-Schatz stent, the '984 patent and the connected z-stent art).
- Trial Transcript from Nov. 8, 2000 at 547-63, 657-63, 674-722, 782-85 (Cordis expert testimony regarding the Palmaz-Schatz stent, the '984 patent and the connected z-stent art).
- Trial Transcript from Nov. 9, 2000 at 819-23, 921 (Cordis expert testimony regarding the '984 patent); 926-941 (R. Croc testimony re Palmaz-Schatz stent); 1033-1053 (R. Schatz testimony).
- Trial Transcript from Nov. 13, 2000 at 1086-1134 (R. Schatz testimony); 1275-1305 (Cordis expert testimony regarding the '984 patent).
- Trial Transcript from Nov. 14, 2000 at 1390-1404, 1448-1454, 1486-1500 (Cordis expert testimony regarding the '984 patent).
- Trial Transcript from Nov. 15, 2000 at 1686-87, 1724-42, 1828-34, 1850-54, 1887-92 (AVE expert testimony regarding the '984 patent).
- Trial Transcript from Nov. 16, 2000 at 2077-198 (AVE expert testimony regarding the alleged obviousness of the '984 patent).
- Trial Transcript from Nov. 17, 2000 at 2331-34 (jury instructions as to the meaning of the limitations of the claims of the '984 patent).
- Trial Transcript from Nov. 20, 2000 at 2441-48, 2499-2500, 2546-50, 2552-56 (Attorney's closing arguments regarding the '984 patent).
- Trial Transcript from Nov. 21, 2000 at 2592-94 (reading of jury verdict).
- Trial Transcript from Dec. 18, 2000 at 2750-95 (Cordis expert testimony regarding the Palmaz-Schatz stent during the damages phase).
- Trial Transcript from Dec. 20, 2000 at 3421-88 (AVE expert testimony regarding the Palmaz-Schatz stent during the damages phase).
- Jury verdict, dated Nov. 21, 2000.
- District Court decisions on post-trial motions (194 F. Supp. 2d 323).
- Court of Appeal for the Federal Circuit decision (339 F.3d 1352).

A1239

## US 7,300,662 B2

Page 7

Trial Transcript from Mar. 4, 2005 at 133-135, 171-173 and 192-96 (Attorney's opening remarks regarding '984 validity).

Trial Transcript from Mar. 7, 2005 at 275-311 (Cordis expert testimony regarding the Palmaz-Schatz stent); 342-46, 353-59, 416-425 (J. Palmaz testimony regarding the Palmaz-Schatz stent, the '984 patent and the connected z-stent art); 430-449, 452-58, 462-492 (R. Croce testimony regarding the Palmaz-Schatz stent); 500-507 (Cordis expert testimony regarding the '984 patent).

Trial Transcript from Mar. 8, 2005 at 609 (Cordis expert testimony regarding the '984 patent); 628-73, 724-740, 773, 801-839 (Cordis expert testimony regarding the '984 patent, the prior art and the Palmaz-Schatz stent).

Trial Transcript from Mar. 9, 2005 at 936-49, 968-69 (Cordis expert testimony regarding the '984 patent, the prior art and the Palmaz-Schatz stent).

Trial Transcript from Mar. 10, 2005 at 1427-74, 178-1509, 1514-23 (AVE expert testimony regarding the alleged obviousness of the '984 patent); 1566-93 (AVE expert testimony regarding Palmaz-Schatz stent); 1634-49 (R. Schatz testimony).

Trial Transcript from Mar. 11, 2005 at 1846-47, 1891-1900, 1919 (Attorneys' closing arguments regarding '984 obviousness).

Trial Transcript from Mar. 14, 2005 at 1964-67 (reading of jury verdict).

Jury verdict dated Mar. 14, 2005.

Medtronic Vascular Inc.'s Opening Brief in Support of Its Motion for Judgment As A Infringement Claim dated Apr. 19, 2005.

Medtronic Vascular Inc.'s Opening Brief in Support of Its Motion for a New Trial dated Apr. 19, 2005.

D.I. 1407, Cordis' Combined Answering Brief In Opposition to AVE's Motion for JMOL on Infringement of the Palmaz '762 and Schatz '984 Patents and Its Motion for a New Trial dated May 5, 2005.

D.I. 1414, Medtronic Vascular Inc.'s Combined Reply Brief In Support of Its Motion for Judgment as a Matter of Law on Cordis Corp.'s Patent Infringement Claims and Its Motion for a New Trial dated May 19, 2005.

Trial Transcript from Feb. 8, 2001 at 372-412, 449-469 (B. Tobor testimony regarding the prosecution of the '417, '984 and '332 patents); 510-13 (J. Milnamow testimony regarding the prosecution of the '332 patent); 558-604 (J. Palmaz testimony regarding the prosecution of the '417, '984 and '332 patents and the prior art).

Trial Transcript from Feb. 9, 2001 at 637-45, 662-672, 682-85 (J. Palmaz testimony regarding the prior art); 699-742 (R. Schatz testimony); 769-770, 790-95 (Cordis expert testimony regarding prior art).

D.I. 1067, Medtronic AVE, Inc.'s Post-Trial Brief Relating to the Unenforceability of the '762 and '984 Patents Due to Inequitable Conduct.

D.I. 1077, Cordis' Combined Answering Brief in Opposition to AVE's BSC's Post-Hearing Briefs on Alleged Inequitable Conduct Concerning the '762, '984 and '332 Patents.

D.I. 1089, Reply Brief In Support of Medtronic AVE, Inc.'s Contention that the '762 and '984 Patents are Unenforceable Due to Inequitable Conduct dated May 7, 2001.

C.A. No. 00-886-SLR, Answer and Counterclaims of Def. Medtronic AVE, Inc. To First Amended Complaint of Plaintiff Cordis Corp.

BSC's Opening Post-Trial Brief in Support of Its Defense That the Patents in Suit Are Unenforceable, dated Mar. 16, 2001.

Reply Brief in Support of BSC's Defense That the Patents in Suit Are Unenforceable, dated May 7, 2001.

Court's Decision on allegations of inequitable conduct (194 F. Supp. 2d 323) Mar. 28, 2002.

Trial Transcript from Nov. 21, 2000 at 155-57 and 180-84 (Attorneys' opening remarks regarding '332 patent).

Trial Transcript from Nov. 27, 2000 at 227-51, 260-300 (Cordis expert testimony regarding the Palmaz-Schatz stent); 343-60, 363-67, 424-33 (J. Palmaz testimony regarding the Palmaz-Schatz stent and the '332 patent).

Trial Transcript from Nov. 28, 2000 at 649-71.

Trial Transcript from Nov. 29, 2000 at 791-816, 859-870, 953-62 (Cordis expert testimony regarding the '332 patent and the Palmaz-Schatz stent).

Trial Transcript from Nov. 30, 2000 at 1018 (Cordis expert testimony regarding the '332 patent); 1062-80, 1108-1111 (R. Croce testimony regarding the Palmaz-Schatz stent); 1169-70, 1205-17, 1236-45 (Cordis expert testimony regarding the '332 patent).

Trial Transcript from Dec. 1, 2000 at 1352-54 (Cordis expert testimony regarding the '332 patent); 1364-1442 (R. Schatz testimony); 1493-1508, 1552-69 (BSC expert testimony regarding the '332 patent and the Palmaz-Schatz stent).

Trial Transcript from Dec. 4, 2000 at 1602-12, 1638-51, 1713-14, 1730-61, 1811-14, 1823-36 (BSC expert testimony regarding the alleged obviousness of the '332 patent, the prior art and the Palmaz-Schatz stent).

Trial Transcript from Dec. 6, 2000 at 2318-27, 2342-58 (BSC expert testimony regarding the '332 patent).

Trial Transcript from Dec. 7, 2000 at 2549-52 (Cordis expert testimony regarding the '332 patent); 2575-2579, 2591-92, 2630-31, 2649, 2669-71, 2684-85, 2688, 2708-10, 2725-27 (Attorney closing argument regarding '332 patent); 2742-46 Q'ury instructions as to the meaning of the limitations fo the claims of the '332 patent).

Trial Transcript from Dec. 11, 2000 at 2817-22 (reading of jury verdict).

Jury verdict, dated Dec. 11, 2000.

D.I. 699, Motion by Defendant BSC and Scimed Life Systems, Inc. For Summary Judgment of Invalidity of U.S. Patent No. 5,902,332 dated Apr. 4, 2000.

D.I.896, Order Denying Motion for Summary Judgment of Invalidity and Unenforceability of Claims 1, 3, and 5 of the U.S. Patent No. 5,902,332 Denying {699-1} Motion for Summary Judgment of Invalidity of U.S. Patent No. 5,902,332 dated Oct. 12, 2000.

Wright et al., Percutaneous Endovascular Stent: An Experimental Study (Abstract), RSNA Meeting (Nov. 28, 1984).

Hearing Transcript from Feb. 10, 1998 at 122-32, 146-80 (Attorneys' opening remarks regarding '417 patent); 180-312 (R. Schatz testimony) [Portions of This Transcript Have Been Removed As Confidential].

Hearing Transcript from Feb. 11, 1998 at 427-575, 577-651 (Cordis expert testimony regarding the '417 patent, the prior art and the Palmaz-Schatz stent).

Hearing Transcript from Feb. 13, 1998 at 1121-1261 (Guidant expert testimony regarding the alleged obviousness of the '417 patent, the prior art and the Palmaz-Schatz stent). [Portions of This Transcript Have Been Removed As Confidential].

Order by J. Robinson denying Cordis' Motion for a Preliminary Injunction Against ACS dated Jul. 17, 1998.

ACS, Inc.'s and Guidant Corp.'s Opening Brief in Support of Their Motion for Summary Judgment of Invalidity of U.S. Patent No. 5,102,417 dated Aug. 27, 1998.

Plaintiffs' Answering Brief in Opposition to ACS' and BSC' Motion for Summary Judgment on Obviousness dated Sep. 24, 1998.

Order dated Mar. 31, 2000.

Schatz Deposition Testimony; May 15, 1996: 79-83, 89-92, 105-107 and 153-161.

Schatz Deposition Testimony; May 16, 1996: 555-564, 569-572.

Schatz Deposition Testimony; Jan. 18, 1998: 67-73, 108-110.

Schatz Deposition Testimony; Jul. 14, 1998: 69-77, 108-112, 119-123.

Schatz Deposition Testimony; Jul. 12, 1999: 88-91, 132-135, 144-149, 218-223, 231-242.

Schatz Deposition Testimony; Jul. 13, 1999: 251-334, 339-345, 374-416.

Schatz Deposition Testimony; Jul. 14, 1999: 454-550.

Schatz Deposition Testimony; Jul. 15, 1999: 560-614.

Schatz Deposition Testimony; Dec. 2, 1999: 906-911, 928-942, 945-963, 976-978, 1029-1034, 1038-1042.

Palmaz Deposition Testimony, Nov. 5, 1991: 160-172.

Palmaz Deposition Testimony, Feb. 5, 1995: 710-727.

Palmaz Deposition Testimony, Jul. 16, 1998: 55-56; 81-82.

Palmaz Deposition Testimony, Jul. 28, 1999: 560-568, 570-579.

Palmaz Deposition Testimony, Jul. 29, 1999: 778-785.

Palmaz Deposition Testimony, Aug. 31, 1999: 1403-1452.

Palmaz Deposition Testimony, Sep. 2, 1999: 1953-1960.

## US 7,300,662 B2

Page 8

- Palmaz Deposition Testimony, Oct. 14, 1999: 2201-2209; 2275-2342; 2371-2411.
- Palmaz Deposition Testimony, Oct. 15, 1999: 2424-2497; 2508-2589.
- Palmaz Deposition Testimony, Oct. 16, 1999: 2853-2860.
- Tobor Deposition Testimony, Jun. 17, 1999: 837-958.
- Tobor Deposition Testimony, Jun. 18, 1999: 1095-1184.
- Tobor Deposition Testimony, Dec. 1, 1999: 1217-1371.
- Tobor Deposition Testimony, Dec. 2, 1999: 1398-1414; 1444-1508; 1532-1548.
- Tobor Deposition Testimony, Dec. 3, 1999: 1652-1653; 1662-1672; 1683-1694.
- Kula Deposition Testimony, Apr. 20, 1999: 268-169.
- Kula Deposition Testimony, Nov. 16, 1999: 660-675; 680-694; 748-755; 774-821.
- Kula Deposition Testimony, Nov. 18, 1999: 176-223.
- Expert Report of Dr. Rodney S. Badger on Behalf of Medtronic AVE, Inc. (Jan. 31, 2000).
- Expert Report of Dr. Joseph Bonn on Behalf of Medtronic AVE, Inc. (Jan. 31, 2000).
- Deposition of Dr. Joseph Bonn dated Mar. 14, 2000.
- Rebuttal Expert Report of Nigel Buller, B.Sc., M.D., F.R.C.P. (Mar. 2000).
- Second Supplemental Rebuttal Expert Report of Nigel Buller, B.Sc., M.B., F.R.C.P. (Aug. 17, 2004).
- Rebuttal Expert Report of John M. Collins, PH.D. (Feb. 2000).
- Expert Report of David C. Cumberland, M.D. (Jan. 24, 2000).
- Expert Report of John T. Goolkasian (Feb. 2000).
- Deposition of Richard R. Heuser, M.D. (Sep. 7, 2004).
- Deposition of Henry R. Piehler (Sep. 10, 2004).
- Deposition of Ronald J. Solar (Mar. 22, 2000).
- Deposition of Ronald J. Solar (Mar. 23, 2000).
- Deposition of Ronald J. Solar (Apr. 12, 2000).
- Expert Report of Dr. Arina Van Breda on Behalf of Medtronic AVE, Inc. (Jan. 31, 2000).
- Deposition of Arina Van Breda (Mar. 24, 2000).
- Deposition of Arina Van Breda (Aug. 21, 2004).
- Expert Report of John F. Witherspoon (Jan. 24, 2000).
- Supplemental Expert Report of John F. Witherspoon (Oct. 27, 2000).
- Deposition of John F. Witherspoon (Mar. 8, 2000).
- Palmaz et al., Article: Normal and Stenotic Renal Arteries: Experimental Balloon Expandable Intraluminal Stenting, *Radiology*, Sep. 1987. (AVE 84).
- Julio C. Palmaz, Article: "Expandable vascular endoprosthesis." (AVE 132).
- Duprat et al., Article: Flexible Balloon-Expandable Stent for Small Vessels Duprat et al. *Radiology*, vol. 162, pp. 276-278, 1987. (AVE 134).
- Coons et al., Article: "Large-Bore, Long Biliary Endoprosthesis (Biliary Stents) for Improved Drainage," *Radiology*, vol. 148, pp. 89-94, 1983. (AVE 143).
- Honickman et al., Article: "Malpositioned Biliary Endoprosthesis, Technical Developments And Instrumentation," vol. 144, No. 2, 1982. (AVE 144).
- Harries-Jones, et al., Article: "Repositioning of Biliary Endoprosthesis with Gruntzig Balloon Catheters," *AJR*, vol. 138, pp. 771-772, 1982. (AVE 153).
- Charnsangavej et al., Article "Stenosis of the Vena Cava: Preliminary Assessment of Treatment with Expandable Metallic Stents," *Radiology*, vol. 161, pp. 295-298, 1986. (AVE 359).
- Article "Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications," *Radiology*, vol. 158, pp. 309-312, 1986. (AVE 364).
- T. Yoshioka, et al., *AJR* Article: "Self-Expanding Endovascular Graft: An Experimental Study in Dogs", vol. 151, pp. 673-676, 1988. (AVE 438).
- Article: "Expandable Intraluminal Vascular Graft: A Feasibility Study," *Surgery*, vol. 99, pp. 199-205, 1986. (AVE 461).
- Lawrence et al., Article: "Percutaneous Endovascular Graft: Experimental Evaluation." *Radiology*, vol. 163, pp. 357-360, 1987. (AVE 671).
- Palmaz et al., Article: Expandable Intraluminal Graft: A Preliminary Study, Nov. 17-22, 1985, *Radiology*, vol. 156, pp. 73-77, 1985. (AVE 1224).
- Fallone et al., "Elastic Characteristics of the Self-Expanding Metallic Stents," *Investigative Radiology*, vol. 23, pp. 370-376, 1988. (AVE 1953).
- Palmaz Paper Entitled "Research Project Expandable Vascular Endoprosthesis" May 18, 1983.
- Rousseau, et al., Publication: "Percutaneous Vascular Stent: Experimental Studies & Preliminary Clinical Results in Peripheral Arterial Diseases," in *Inter. Angio*, vol. 6, 153-161, 1987. (AVE 3301).
- Rousseau, et al., Publication: "Self-Expanding Endovascular Prosthesis: An Experimental Study," *Radiology*, vol. 164, pp. 709-714, 1987. (AVE 3303).
- Wallace, et al., Article: "Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications," *Radiology*, vol. 58, pp. 309-312, 1986. (DBX 2938).
- Palmaz et al., Article: "Expandable Intraluminal Graft: A Preliminary Study," *Radiology*, vol. 156, pp. 73-77 (DBX 4595).
- Program for the 12th Annual Course on Diagnostic Angiography and Interventional Radiology Mar. 23-26, 1987 sponsored by The Society of Cardiovascular and Interventional Radiology (DBX 6235).
- Preliminary Motion for Judgment re: Wolff claims 1, 2-8, 10, 15 and 19 (DBX6759).
- Palmaz Declaration (DBX 7069).
- Letter from Gaterud to Dr. Palmaz dated Jul. 5, 1988 with attached document entitled: "Segmented, balloon-expandable stents." (DBX 7160).
- Duprat et al., Article: "Flexible Balloon-Expandable Stent For Small Vessels," *Radiology*, vol. 168, pp. 276-278, 1987 (PX 82).
- Drawing Sent to Bodic on Mar. 17, 1986 (PX 374).
- Letter from Dr. Palmaz to R. Bowman enclosing a model of the flexible coronary graft dated Mar. 17, 1986 (PX 337).
- Lab Notebook pages dated Jul. 30, 1987 from Rodney Wolff (COR 185596-597) (PX621A).
- Charnsangavej, et al., Article: "Stenosis of The Vena Cava Preliminary Assessment of Treatment with expandable Metallic Stents," *Radiology*, vol. 161, No. 2, pp. 295-298 with attached photographs, 1986. (API 72).
- J. Palmaz: The Current Status of Vascular Prostheses, published by SCIR in the Twelfth Annual Course on Diagnostic Angiography And Interventional Radiology Mar. 23-26, 1987. (API 73).
- Amendment in Response to Office Action of Oct. 18, 1988 in re: Application of Julio Palmaz S/N 174,246. (API 152).
- Article: Wallace, et al., Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications Work In Progress, *Radiology*, vol. 158, pp. 309-312. (API 295).
- Reply of Senior Party Schatz To Patentee Wolff's Opposition To The Belated Motion For Judgment Of Applicant Schatz With Regard To Wolff Claims 1, 2-8, 10, 11, 13-17, And 19 (COR 186450-455) (API 310).
- Brief Of Senior Party Schatz At Final Hearing (API 313).
- Letter from Ron Sickles to Ben Tobor dated Feb. 10, 1988 (Exhibit 42).
- Letter from R.O. Sickles to Mike Tatlow dated May 12, 1988 (Exhibit 43).
- Letter from R. O. Sickles to Richard Schatz dated Jun. 2, 1988 (Exhibit 44).
- Letter from Richard Schatz to Raimund Erbel dated Jun. 3, 1988 (Exhibit 45).
- Letter from Richard Schatz to Mike Schuler dated Aug. 29, 1991 (Exhibit 48).
- Minutes of J&J Stent Project Review Meeting dated Jan. 21, 1988 (Exhibit 7).
- Preliminary Motion for Judgment with Regard to Wolff Claims 1, 2-8, 10, 11, 13-17, and 19. (Exhibit 67).
- Declaration of Richard A Schatz. (Exhibit 75).
- Belated Motion for Judgement with Regard to Wolff Claims 1, 2-8, 10, 11, 13-17 and 19. (Schatz—Exhibit 77).
- Letter from Dr. Schatz to Mr. Tobor, dated Jun. 3, 1988. (Exhibit 122).

A1241

## US 7,300,662 B2

Page 9

Letter from Dr. Schatz to Mr. Romano, dated Nov. 28, 1988. (Exhibit 131).

Letter from Mr. Sickles to Mr. Tobor, dated Feb. 10, 1988 (Exhibit 145).

Richard A. Schatz, Article titled: "A View of Vascular Stents" *Circulation*, vol. 79, No. 2, pp. 445-457, 1989. (Exhibit 194).

Senior Party Schatz's reply to Patente Wolffs Opposition to the Preliminary Motion Of Applicant Schatz for Judgment with regard to Wolff Claims 1, 2-8, 10, 11, and 13-17. (Exhibit 69).

Wallace, et al., Article: "Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications' Work In Progress," *Radiology*, vol. 158, pp. 309-312, 1986. (Exhibit 165).

Charnsangavej, et al., Article: "Stenosis of The Vena Cava Preliminary Assessment of Treatment with expandable Metallic Stents," *Radiology*, vol. 161, No. 2, pp. 295-298 with attached photographs, 1986. (Exhibit 167).

David D. Lawrence et al., Publication: Percutaneous Endovascular Graft: Experimental Evaluation<sup>1</sup>, *Radiology*, pp. 357-360, 1987. (Exhibit 173).

Charles E. Putnam, M.D., Cover and article from "Investigative Radiology", vol. 23, No. 5, May 1988. (Exhibit 177).

Robert N. Berk, Cover and article from "American Journal of Roentology", pp. 673-676, 1988. (Exhibit 178).

Declaration of John S. Kula Under 37 CFR § 1.672. (Kula—Exhibit 77).

Yoshioka et al., Article: "Self-Expanding Endovascular Graft: An Experimental Study in Dogs" *AJR*, vol. 151, pp. 673-676, 1988. (PX 100).

Palmaz, et al., Article: Expandable Intraluminal Graft: A Preliminary Study Work in Progress<sup>1</sup>, *Radiology*, vol. 156, No. 1, pp. 73-77, 1985. (PX 101).

Declaration of Richard Schatz Under 37 C.F.R. § 1.672. (PX 106).

Charnsangavej et al., Article: "Stenosis of the Vena Cave: Preliminary Assessment of Treatment with Expandable Metallic Stents," *Radiology*, vol. 161, pp. 295-298, 1986. (PX 143).

Wallace, et al., Article: Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications Work in Progress<sup>1</sup>, *Radiology*, vol. 158, pp. 309-312, 1986. (PX 144).

Gina Kolata, News Article: NY Times, "Devices That Opens Clogged Arteries Gets a Falling Grade in a New Study", pp. 16-18, Jan. 3, 1991. (PX 186).

Duprat, et al., Article: "Flexible Balloon- Expanded Stent for Small Vessels Work in Progress"<sup>1</sup>, *Radiology*, vol. 162, pp. 276-278, 1987. (PX 207).

Letter from Palmaz to Bowman dated Mar. 17, 1986. (PX 350).

Memo re: Minutes of Stent Project Review- San Antonio- Mar. 15, 1988. (PX 651).

Kuntz, et al., Article: Clinical Cardiology Frontiers: "Defining Coronary Restenosis, Newer Clinical and Angiographic Paradigms", *Circulation*, Sep. 1993, vol. 88, No. 3, pp. 1310-1323. (PX 854).

Belated Motion for Judgment with regard to Wolff Claims<sup>1</sup>, 2-8, 10, 11, 13-17, and 19. (PX 1410).

Drawing of Sprial Stent (sent to Bodic Mar. 17, 1986). (PX2933).

Wright et al., Article: "Percutaneous Endovascular Stents: An Experimental Evaluation," *Radiology*, vol. 156, pp. 69-72, 1985. (PX 3093).

Charnsangavej et al., Article: "A New Expandable Metallic Stent for Dilation of Stenotic Tubular Structures: Experimental and Clinical Evaluation," *Houston Medical Journal*, vol. 3, pp. 41-51, Jun. 1987. (PX 3207).

In re Application of Wiktor, Appln. No. 69,636, Response to Office Action dated Mar. 17, 1988. (PX3236).

Transmittal Letter of Response to First Office Action in '417 patent. (PX 3993).

Letter from B. Tobor to R. Schatz dated Jul. 23, 1991. (PX 3996).

Mullins et al., Article: "Implantation of balloon-expandable intravascular grafts by catheterization in pulmonary arteries and systemic veins," *Circulation*, vol. 77, No. 1, pp. 188-189, 1988. (PX4049).

Schatz et al., Article: "Intravascular Stents for Angioplasty," *Cardio*, 1997. (PX 4050).

Schatz et al., Article: "New Technology in Angioplasty Balloon-Expandable Intravascular Stents, New Developments in Medicine," vol. 2, No. 2 pp. 59-75, 1987. (PX4051).

Richard A. Schatz, Article: "Introduction to Intravascular Stents," *Cardiology Clinics*, vol. 6, No. 3, pp. 357-372, 1988. (PX 4052).

Richard A. Schatz, Article: "A View of Vascular Stents," *Circulation*, vol. 79, No. 2, pp. 445-457, 1989. (PX4053).

Wang et al., Article: "An Update on Coronary Stents," *Cardio*, pp. 177-186, 1992. (PX 4054).

Richard A. Schatz, Article: "New Technology in Angioplasty: Balloon-Expandable Starts," *Medicamundi*, vol. 33, No. 3, pp. 112-116, 1988. (PX 4055).

Letter from Tobor to Schatz dated Sep. 29, 1988. (PX 1395).

Verified Statement of Facts by Unnamed Inventor R.A. Schatz document filed in U.S. Patent and Trademark Office on Sep. 8, 1989. (PX 3677).

Declaration of John S. Kula Under 37 CFR § 1.672 (Exhibit 329).

Letter to Mike Schular from R.A. Schatz dated Aug. 29, 1991. (Exhibit 402).

Articulated, Balloon—Expandable Stents, (DBX 7159).

J. Rosch et al., Experimental Intrahepatic Portacaval Anastomosis: Use of Expandable Gianturco Stents, *Radiology*, vol. 162, pp. 481-485, 1987.

J. Rosch et al., Modified Gianturco Expandable Wire Stents In Experimental and Clinical Use, *Ann Radiol*, vol. 31, No. 2, pp. 100-103, 1987.

J. Rosch et al., Gianturco Expandable Stents In the Treatment of Superior Vena Cava Syndrome Recurring After Vena Cava Syndrome Recurring After Maximum-Tolerance Radiation, *Cancer*, vol. 60, pp. 1243-1246, 1987.

I.E. Gordon, Structures or Why Things Don't Fall Down, Penguin Books, pp. 45-59, 132-148, 210-244, 377-383.

Maass et al., Radiological Follow-up of Transluminally Inserted Vascular Endoprostheses: An Experimental Study Using Expanding Spirals, *Radiology*, vol. 152, pp. 659-663, 1984.

Argument submitted re EP 861 15473 dated Jan. 20, 1995. (AVE 2478).

Verified Statement of Facts by Julio C. Palmaz dated Aug. 4, 1989. (PX 3662).

Papanicolaou et al., Insertion of a Biliary Endoprosthesis Using A Balloon Dilatation Catheter, *Gastrointest Radiology*, vol. 10, pp. 394-396, 1985.

Palmaz et al., Atherosclerotic Rabbit Aortas: Expandable Intraluminal Grafting, *Radiology*, vol. 168, pp. 723-726, 1986.

Palmaz, The Current Status of Vascular Prostheses; Rosch et al., Gianturco, Expandable Stents in Experimental and Clinical Use, *SCIVR*, pp. 118-124, 1987.

Rosch et al., Abstract: Modified Gianturco Expandable Wire Stents in Experimental and Clinical Use, *CIRSE, Porto Cervo, Sardinia*, May 25-29, 1987.

Rosch et al., Gianturco Expandable Wire Stents in the Treatment of Superior Vena Cava Syndrome Recurring After Maximum-Tolerance Radiation, *Cancer*, vol. 60, pp. 1243-1246, 1987.

Mirich et al., Percutaneously Placed Endovascular Grafts for Aortic Aneurysms: Feasibility Study, *Radiology*, vol. 170, pp. 1033-1037, 1989.

Dotter, Transluminally-placed Coilspring Endarterial Tube Grafts, *Investigative Radiology*, vol. 4, Sept.-Oct., pp. 329-332, 1969.

Palmaz et al., Abstract: Expandable Intraluminal Graft: A Preliminary Study, *Radiology*, vol. 153 (P), Nov. 1983: 70<sup>th</sup> Scientific Assembly and Annual Meeting.

Cragg et al., Nonsurgical Placement of Arterial Endoprostheses: A New Technique Using Nitinol Wire, *Radiology*, vol. 147, pp. 261-263, Apr. 1983.

J. Rosch et al., Gianturco Expandable Stents in Experimental and Clinical Use, Program: "Twelfth Annual Course on Dignostic Angiography and Interventional Radiology" (Society of Cardiovascular and Interventional Radiology, Pittsburgh, PA), Mar. 23-26, 1987 (the second Monofilament Article).

Uchida et al., Modifications of Gianturco Expandable Wire Stents, *AIR*, vol. 150, pp. 1185-1187, 1988.

Palmaz, Balloon-Expandable Intravascular Stent, *AJR*, vol. 1510, pp. 1263-1269.

## US 7,300,662 B2

Page 10

*Cordis Corporation v. Advanced Cardiovascular Systems, Inc., Guidant Corporation, Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCMED Life System, Inc.*, Plaintiffs Complaint, Oct. 23, 1997 (Case No. 97-550-SLR).

*Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership*, Plaintiffs First Amended Complaint for Declaratory Relief of Patent Validity, Unenforceability, Noninfringement, and for Antitrust Violations, Jan. 27, 1998 (Civil Action No. 97-700).

*Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Cordis Corporation and Johnson & Johnson's Answer and Counterclaim*, Feb. 27, 1998 (Civil Action No. 97-700-SLR).

*Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Expandable-Graft Partnership's Answer*, Feb. 27, 1998 (Civil Action No. 97-700-SLR).

*Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership*, Reply of Plaintiff Arterial Vascular Engineering, Inc. To Counterclaims of Defendant Cordis Corporation, Mar. 31, 1998 (Civil Action No. 97-700-SLR).

*Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership*, Reply of Plaintiff Arterial Vascular Engineering, Inc. To Counterclaims of Defendant Expandable Grafts Partnership, Mar. 31, 1998 (Civil Action No. 97-700-SLR).

*Cordis Corporation v. Advanced Cardiovascular Systems, Inc. and Guidant Corporation* Cordis Corporation's Motion for a Preliminary Injunction, Oct. 8, 1997 (Civil Action No. 97-550).

*Cordis Corporation v. Advanced Cardiovascular Systems, Inc., Guidant Corporation Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCJVTED, Inc.*, *Cordis's Motion for Preliminary Injunction Against Arterial Vascular Engineering, Inc.*, Dec. 29, 1997 (Case No. 97-550-SLR).

Deposition of R. Schatz, M.D. in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, taken on Jan. 8, 1998 (Civil Action No. 97-550 SLR).

Deposition of Lee P. Bendel in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, taken on Jan. 22, 1998 (Civil Action No. 97-550 SLR).

Deposition of Julio Cesar Palmaz in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, taken on Dec. 29, 1997 (Civil Action No. 97-550 SLR).

Deposition of Richard A. Bowman in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, taken on Jan. 9, 1998 (Civil Action No. 97-550 SLR).

Deposition of Gary Schneiderman in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, taken on Jan. 16, 1998 (Civil Action No. 97-550 SLR).

Deposition of David Pearle, M.D. in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, taken on Jul. 10, 1998 (Civil Action No. 97-550 SLR).

Preliminary Injunction hearing testimony taken on Feb. 9-13, 1998 (Civil Action No. 97-550 SLR).

*Cordis Corporation v. Advanced Cardiovascular Systems, Inc.*, et al., (Civil Action No. 97-550 SLR) and *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.* Et al. (Civil Action No. 98-65-SLR), Opening Post Hearing Brief of Plaintiff Cordis Corporation in Support of Motion for Preliminary Injunction, Mar. 6, 1998 (Portions relevant to patent claim construction and patent validity issues).

*Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc.* et al., Post-Hearing Reply Brief of Plaintiff Cordis Corporation in Support of Its Motion for Preliminary Injunction, Apr. 10, 1998 (Case No. 97-550 SLR) (Portions relevant to patent validity issues).

*Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc.*, et al., Plaintiffs Motion for a Preliminary Injunction Against Boston Scientific Corporation and SCLMED Life Systems, Inc. And Memorandum in Support, Apr. 13, 1998 (Case No. 97-550-SLR).

*Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc.*, et al., Judge Robinson's

Order Denying Plaintiffs Motion for a Preliminary Injunction, Jul. 17, 1998 (Civil Action No. 97-550 SLR).

*Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc.*, et al., Defendant Boston Scientific Corporation and SCTMED Life Systems, Inc.'s Motion for Summary Judgment of Invalidity of U.S. Patent No. 5,102,417, Aug. 27, 1998 (Civil Action No. 97-550-SLR).

*Boston Scientific Limited*, et al. v. *Expandable Grafts Partnership*, Plaintiff's Statement of Claim, Mar. 13, 1997 (UK Action No. 1493).

*Boston Scientific Limited*, et al. v. *Expandable Grafts Partnership*, Defendant's Amended Defense and Counterclaim, Aug. 14, 1997 (UK Action No. 1493).

*Boston Scientific Limited*, et al. v. *Expandable Grafts Partnership*, Petition for Revocation, Mar. 13, 1997 (UK Action No. 1497).

*Boston Scientific Limited*, et al. v. *Expandable Grafts Partnership*, Particulars of Objections, Mar. 13, 1997 (UK Action No. 1497).

*Boston Scientific Limited*, et al. v. *Expandable Grafts Partnership and Boston Scientific Limited* et al., v. *Julio C. Palmaz, Boston's Skeleton Argument* (UK Action Nos. 1493, 1495, 1496, and 1497). *Boston Scientific Limited*, et al. v. *Julio C. Palmaz and Expandable Grafts Partnership*, Skeleton Argument of Palmaz/EGP, Mar. 19, 1998 (UK Action Nos. 1493, 1495, 1496 and 1497).

*Boston Scientific Limited*, et al. v. *Julio C. Palmaz and Expandable Grafts Partnership*, EGP's Final Submissions, Apr. 2, 1998 (UK Action Nos. 1493, 1495, 1496 and 1497).

*Boston Scientific Limited*, et al. v. *Julio C. Palmaz and Expandable Grafts Partnership*, Judgment, Jun. 26, 1998 (UK Action Nos. 1493, 1495, 1496 and 1497).

Rosch, Modified Gianturco Expandable Wire Stents in Experimental and Clinical Use, CJJR.SE 1987 Presentation: see Witness Statement of Josef Rosch from U.K. Proceeding.

Statement of Claim by Boston Scientific et al. against Expandable Grafts Partnership et al., in *EPG* et al., v. *Boston Scientific* et al. in Netherlands (Mar. 13, 1997).

Motion for Joinder of Actions, Change of Claim and Statement of Claim filed by Expandable Grafts Partnership et al. in *EPG* et al. v. *Boston Scientific* et al. In Netherlands (Apr. 22, 1997).

Opinion of K.J. Merman filed in *EPG* et al. v. *Boston Scientific* et al. in Netherlands (Aug. 29, 1997).

Expert report of Dr. Nigel Buller in *EPG* et al. v. *Boston Scientific* et al. in Netherlands (Aug. 28, 1997).

Expert report of Lee P. Bendel in *EPG* et al. v. *Boston Scientific* et al. in Netherlands (Aug. 28, 1997).

Memorandum or Oral Pleading in *EPG* et al. v. *Boston Scientific* et al. in Netherlands (Sep. 12, 1997).

Plea Notes of P. A.M. in *EPG* et al. v. *Boston Scientific* et al. in Netherlands (Mar. 10, 1998).

Decision of Court of Appeals in *EPG* et al. v. *Boston Scientific* et al. In Netherlands (Apr. 23, 1998).

Translation of Nullity Action Against EPO 0 364 787 by Biotronik in Germany.

Translation of Nullity Action Against EPO 0 335 341 by Biotronik in Germany.

Translation of EPG Response to Nullity Action Against EP 0 364 787 by Biotronik in Germany.

Translation of EPG Response to Nullity Action EP 0 335 341 by Biotronik in Germany.

Nullity Suit Against EP-B1-0 335 341 Brought by Boston Scientific in Germany.

Translation of Opposition filed by Terumo Corp. Against Japan Patent No. 2680901.

Translation of Decision on Opposition Against Japan Patent No. 2680901.

Memorandum Order of the Court dated Sep. 7, 2000, concerning disputed claim construction.

Translation of Judgment in Nullity Action Against EP 0 364 787 by Biotronik in Germany.

Translation of Judgment in Nullity Action Against EP 0 335 341 by Biotronik in Germany.

Trial transcript from Mar. 17, 2005 at 171-172, 191-192.

Trial transcript from Mar. 18, 2005 at 282-285, 325-327, 349-351.

Trial transcript from Mar. 21, 2005 at 721-726.



## US 7,300,662 B2

Page 11

Trial transcript from Mar. 24, 2005 at 1387.

Trial transcript from Jul. 26, 2005.

BSC's Opening Brief in Support of Its Motion for Judgment as a Matter of Law or, in the Alternative, for a New Trial, dated Mar. 16, 2001.

Cordis' Answering Brief in Opposition to BSC's Motion for JMOL or a New Trial on the Palmaz '762 Patent and the Schatz '332 Patents, dated Apr. 17, 2001.

BSC's Reply Brief in Support of Its Motion for Judgment as a Matter of Law or, in the Alternative, for a New Trial, dated May 11, 2001.

J. Rosch et al., Abstract, Expandable Gianturco-Type Wire Stents in Experimental Intrahepatic Portacaval Shunts, Program: "72nd Scientific Assembly and Annual Meeting of the Radiological Society of North America", Nov. 30-Dec. 5, 1986, Radiology, vol. 161, pp. 40-41, 1986.

*Cordis Corporation v. Boston Scientific*, Order Dated Mar. 27, 2006 (97-550-SLR).

*Cordis Corporation v. Boston Scientific*, Judgment in a Civil Case Dated Mar. 27, 2006 (97-550-SLR).

*Cordis Corporation v. Boston Scientific*, Memorandum Opinion Dated Mar. 27, 2006 (97-550-SLR).

*Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc., Guidant Corporation, Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCIMED Life Systems, Inc.*, Answer and Counterclaims of Defendant Advanced Cardiovascular Systems, Inc., Apr. 8, 1998 (Case No. 97-550-SLR).

*Boston Scientific Limited et al. v. Expandable Grafts Partnership and Boston Scientific Limited et al. v. Julio C. Palmaz*, Boston's Closing Submissions (UK Action Nos. 1493, 1495, 1496 and 1497).

*Cordis Corporation v. Advanced Cardiovascular Systems, Inc., Guidant Corporation, Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCIMED Life Systems, Inc.*, Defendants' Answer, Nov. 12, 1997 (Case No. 97-550-SLR).

Statement of Rejoinder in the Action on the Merits, Also Including an Amendment of Defendant's Final Position in the Principal Action, as Well as the Provisional Statement of Rejoinder in the Action on the Counterclaim in *EPG et al. v. Boston Scientific et al.* In Netherlands (Feb. 10, 1998).

Statement of Answer in the Ancillary Appeal in *EPG et al. v. Boston Scientific et al.* in Netherlands (Mar. 10, 1998).

Appeal filed by Expandable Grafts Partnership et al. in *EPG et al. v. Boston Scientific et al.* in Netherlands (Nov. 12, 1997).

Title filed by Boston Scientific et al. in *EPG et al. v. Boston Scientific et al.* in Netherlands (Jan. 22, 1998).

Deposition of Richard Schatz, M.D. in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.* taken on Jul. 14, 1998 (Civil Action No. 97-550-SLR).

Jury Verdict form from the *Cordis Corporation et al. v. Boston Scientific Corporation*, et al liability trial, undated.

Trial testimony transcripts from the *Cordis Corporation et al. v. Boston Scientific Corporation et al.* liability trial dated Nov. 27-Dec. 1, Dec. 4-8 and Dec. 11, 2000.

*Boston Scientific SCIMED, Inc. and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.*, Opening Expert Report of Stephen R. Hanson, Ph.D. (Civil Action No. 03-283-SLR).

*Boston Scientific SCIMED, Inc. and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.*, Opening Expert Report of Robson F. Storey, Ph.D. (Civil Action No. 03-283-SLR).

*Boston Scientific SCIMED, Inc. and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.*, Rebuttal Expert Report of Kinam Park, Ph.D. (Civil Action No. 03-283-SLR).

*Cordis Corporation v. Boston Scientific Corporation and SCIMED Life Systems, Inc.* (C.A. No. 03-027-SLR) and *Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.* (C.A. No. 03-283-SLR) Combined Post-Hearing Brief In Support Of Cordis Corporation's Motion For Preliminary Injunction in C.A. No. 03-027-SLR, And In Opposition to Plaintiffs' Motion For Preliminary Injunction in C.A. No. 03-283-SLR.

*Cordis Corporation v. Boston Scientific Corporation and SCIMED Life Systems, Inc.* (C.A. No. 03-027-SLR) *Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.* (C.A. No. 03-283-SLR), Boston Scientific's Opening Post-Hearing Brief.

*Cordis Corporation v. Boston Scientific Corporation and SCIMED Life Systems, Inc.* (C.A. No. 03-027-SLR) *Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.* (C.A. No. 03-283-SLR), Combined Post-Hearing Answering Brief In Support of Cordis Corporation's Motion For Preliminary Injunction In C.A. No. 03-027-SLR, And In Opposition To Plaintiffs Motion For Preliminary Injunction In C.A. No. 03-283-SLR.

Wu et al., Silicone-covered self-expanding metallic stents for the palliation of malignant esophageal obstruction and a review of the literature, *Gastrointestinal Endoscopy*, 1994, pp. 22-33, vol. 40, No. 1, Portland Oregon.

Binmoeller, et al., Silicone-Covered Expandable Metallic Stents in the Esophagus: An Experimental Study, *Endoscopy*, 1992, pp. 416-420, vol. 24, Georg Thieme Verlag Stuttgart New York.

*Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.*, Answering Memorandum in Opposition to Plaintiffs Motion for a Preliminary Injunction and Appendix thereto (Civil Action No. 03-283-SLR).

*Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.*, Plaintiff's Reply Brief in Support of Their Motion for Preliminary Injunction. Rhine, Polymers for Sustained Macromolecule Release: Procedures to Fabricate Reproducible Delivery Systems and Control Release Kinetics, *Journal of Pharmaceutical Sciences*, 1980, pp. 265-270, vol. 69, No. 3.

Langer et al., Controlled Release of Macromolecules From Polymers, *Biomedical Polymers Polymeric Materials and Pharmaceuticals for Biomedical Use*, 1980, pp. 112-137, Academic Press, Inc., New York, NY.

Langer et al., Applications of Polymeric Delivery Systems for Macromolecules and Factors Controlling Release Kinetics.

Rhine et al., A Method to Achieve Zero-Order Release Kinetics From Polymer Matrix Drug Delivery Systems, pp. 67-72.

Langer et al., Polymers for the Sustained Release of Macromolecules: Controlled and Magnetically Modulated Systems, *Better Therapy With Existing Drugs: New Uses and Delivery Systems*, 1981, pp. 179-216, Merck Sharp & Dohme International, Rahway, NJ.

Hsieh, et al., Zero-Order Controlled-Release Polymer Matrices for Micro-and-Macromolecules, *Journal of Pharmaceutical Sciences*, 1983 pp. 17-22, vol. 72, No. 1.

Brown et al., In Vivo and In Vitro Release of Macromolecules from Polymeric Drug Delivery Systems, *Journal of Pharmaceutical Sciences*, 1983, pp. 1181-1185, vol. 72, No. 10.

Langer, Implantable Controlled Release Systems, *Pharmac. Ther.*, 1983, pp. 35-51, vol. 21, printed in Great Britain.

Kost et al., Controlled Release of Bioactive Agents, *Trends in Biotechnology*, 1984, pp. 47-51, vol. 2, No. 2, Elsevier BV Amsterdam.

Bawa et al., An Explanation for the Controlled Release of Macromolecules from Polymers, *Journal of Controlled Release*, 1985, pp. 259-267, vol. 1 Elsevier Science BV Amsterdam.

Leong et al., Polymeric controlled drug delivery, 1987, pp. 199-233, vol. 1/3, Elsevier Science Publishers BV Amsterdam.

Langer, Polymeric Delivery Systems, *Targeting of Drugs 2 Optimization Strategies*, 1989, pp. 165-174, Plenum Press, New York and London.

Langer, Biomaterials in Controlled Drug Delivery; New Perspectives from Biotechnological Advances; *Pharmaceutical Technology*, 1989, pp. 18, 23-24, 26, 28, 30.

Langer, Controlled Release Systems, pp. 115-124.

Laurencin et al., Polymeric Controlled Release Systems: New Methods for Drug Delivery, *Clinics in Laboratory Medicine*, 1987, pp. 301-323, vol. 7, No. 2, WB Saunders Company, Philadelphia.

Langer, Biopolymers in Controlled Release Systems, *Polymeric Biomaterials*, pp. 161-169.

## US 7,300,662 B2

Page 12

- Tsong-Pin Hsu et al., Polymers for the Controlled Release of Macromolecules: Effect of Molecular Weight of Ethylene-vinyl Acetate Copolymer, *Journal of Biomedical Materials Research*, 1985, pp. 445-460, vol. 19.
- Langer, Polymers and Drug Delivery Systems, *Long-Acting Contraceptive Delivery Systems*, 1983, pp. 23-32, Harper & Row, Philadelphia, PA.
- Langer, New Drug Delivery Systems: What the Clinician Can Expect, *Drug Therapy*, 1983, pp. 217-231.
- Langer, et al., Chemical and Physical Structure of Polymers as Carriers for Controlled Release of Bioactive Agents: A Review, *Rev. Macromol. Chem. Phys.*, 1983, pp. 61-126.
- Langer, Polymeric Delivery Systems for Controlled Drug Release, *Chem. Eng. Commun.* 1980, pp. 1-48-vol. 6, Gordon and Breach Science Publishers, Inc. USA.
- Langer, et al., Biocompatibility of Polymeric Delivery Systems for Macromolecules, *Journal of Biomedical Materials Research*, 1981, pp. 267-277, vol. 15.
- Langer, Controlled Release: A New Approach to Drug Delivery, *Technology Review*, 1981, pp. 26-34.
- Langer, et al Sustained Release of Macromolecules from Polymers, *Polymeric Delivery Systems*, PGS. 175-176, Gordon and Breach Science Publishers, New York.
- Langer, Polymers for the Sustained Release of Proteins and other Macromolecules, *Nature*, 1976, pp. 797, 263, 799-800, vol. 263, No. 5580.
- Baker, et al., Controlled Release: Mechanisms and Rates (1974).
- Hanson, et al., In Vivo Evaluation of Artificial Surfaces with a Nonhuman Primate Model of Arterial Thrombosis, *Lab Clin. Med.*, Feb. 1980, pp. 289-304.
- Baker, Controlled Release of Biologically Active Agents (1987) pp. 1-275.
- Cordis Corporation v. Boston Scientific Corporation* (CA. No. 03-27-SLR) and *Boston Scientific Scimed, Inc., v. Cordis Corporation and Johnson & Johnson, Incorporated* (CA. No. 03-283-SLR) Hearing Transcripts for Jul. 21, 2003, Jul. 22, 2003, Jul. 23, 2003.
- Cordis Corporation v. Boston Scientific Corporation* et al. (CA. No. 03-027-SLR), and *Boston Scientific Scimed, Inc. et al. v. Cordis Corporation* et al. (CA. No. 03-283-SLR), Boston Scientific's Post-Hearing Reply Brief and Exhibits Thereto, Sep. 12, 2003.
- Cordis Corporation v. Boston Scientific Corporation* et al. (CA. No. 03-027-SLR), and *Boston Scientific Scimed, Inc. et al. v. Cordis Corporation* et al. (CA. 03-283-SLR), Memorandum Order, Nov. 21, 2003.
- Cordis Corporation v. Boston Scientific Corporation* et al. (CA. No. 03-027-SLR), and *Boston Scientific Scimed, Inc. et al. v. Cordis Corporation* et al. (CA. No. 03-283-SLR), Deposition Transcript of Julio C. Palmaz.
- Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable Grafts Partnership*, Cordis Corporation and Johnson & Johnson's Answer and Counterclaim, Feb. 27, 1998 (Civil Action No. 97-700-SLR).
- Plea Notes in *EPG et al. v. Boston Scientific* et al. in Netherlands (Sep. 12, 1997).
- Provisional Judgment *EPG et al. v. Boston Scientific* et al. in Netherlands (Oct. 29, 1997).
- Trial testimony transcripts from the Cordis Corporation et al. v. Medtronic AVE Inc., et al. liability trial dated Nov. 6-9, 13-17 and 20-21, 2000.
- Jury verdict form from the *Cordis Corporation* et al. v. *Medtronic AVE, Inc.* et al. liability trial.
- Hearing testimony transcript from the consolidated *Cordis Corporation* et al. v. *Medtronic AVE, Inc.* et al. and *Boston Scientific Corporation* et al. inequitable conduct hearing dated Feb. 7-9 and 12, 2001.
- Cordis Corporation v. Medtronic Ave., Inc.* et al. OPINION, 97-550-SLR, dated Mar. 28, 2002.
- Cordis Corporation v. Advanced Cardiovascular Systems, Inc.* et al. (CA. No. 97-550-SLR), *Medtronic AVE, Inc. v. Cordis Corporation* et al. (CA. No. 97-700-SLR), *Boston Scientific Corporation v. Athicon, Inc.* et al. (CA. No. 98-19-SLR), Expert Report of John T. Goolkasian, Esq.
- Cordis Corporation v. Advanced Cardiovascular Systems, Inc.* et al. (CA. No. 97-550-SLR), *Medtronic AVE, Inc. v. Cordis Corporation* et al. (CA. No. 97-700-SLR), *Boston Scientific Corporation v. Athicon, Inc.* et al. (CA. 98-19-SLR), Expert Report of John F. Witherspoon.
- Ruef, Johannes MD, et al.; "Flavopiridol Inhibits Smooth Muscle Cell Proliferation In Vitro and Neointimal Formation In Vivo After Carotid Injury In The Rat"; From the Division of Cardiology and Sealy Center for Molecular Cardiology, University of Texas Medical Branch, Galveston; Accepted Apr. 9, 1999; *Circulation* Aug. 10, 1999; pp. 659-665.
- European Search Report EP 05 25 2466 dated Jul. 26, 2005.
- European Search Report EP 05 25 2631 dated Jul. 26, 2005.
- European Search Report EP 05 25 2478 dated Jul. 26, 2005.
- Schuler, W. et al., "SDZ RAD, A New Rapamycin Derivative: Pharmacological Properties In Vitro and In Vivo," *Transplantation*, Jul. 5, 1997, 64(1), 36-42.

\* cited by examiner


U.S. Patent

Nov. 27, 2007


Sheet 1 of 2

US 7,300,662 B2


**FIG. 1**



**FIG. 2**



**FIG. 3**



US 7,300,662 B2

1

# DRUG/DRUG DELIVERY SYSTEMS FOR THE PREVENTION AND TREATMENT OF VASCULAR DISEASE

## CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. application Ser. No. 09/850,293, filed May 7, 2001, now abandoned, which in turn claims priority of U.S. Provisional Application No. 60/263,979, filed Jan. 25, 2001, U.S. Provisional Application No. 60/263,806, filed January 24, 2001, U.S. Provisional Application No. 60/262,614, filed Jan. 18, 2001, U.S. Provisional Application No. 60/262,461, filed Jan. 18, 2001, and is a continuation-in-part of U.S. Application No. 09/575,480, filed May 19, 2000, now pending, which in turn claims priority of U.S. Provisional Application No. 60/204,417, filed May 12, 2000.

## BACKGROUND OF THE INVENTION

### 1. Field of the Invention

The present invention relates to drugs and drug delivery systems for the prevention and treatment of vascular disease, and more particularly to drugs and drug delivery systems for the prevention and treatment of neointimal hyperplasia.

### 2. Discussion of the Related Art

Many individuals suffer from circulatory disease caused by a progressive blockage of the blood vessels that perfuse the heart and other major organs with nutrients. More severe blockage of blood vessels in such individuals often leads to hypertension, ischemic injury, stroke, or myocardial infarction. Atherosclerotic lesions, which limit or obstruct coronary blood flow, are the major cause of ischemic heart disease. Percutaneous transluminal coronary angioplasty is a medical procedure whose purpose is to increase blood flow through an artery. Percutaneous transluminal coronary angioplasty is the predominant treatment for coronary vessel stenosis. The increasing use of this procedure is attributable to its relatively high success rate and its minimal invasiveness compared with coronary bypass surgery. A limitation associated with percutaneous transluminal coronary angioplasty is the abrupt closure of the vessel which may occur immediately after the procedure and restenosis which occurs gradually following the procedure. Additionally, restenosis is a chronic problem in patients who have undergone saphenous vein bypass grafting. The mechanism of acute occlusion appears to involve several factors and may result from vascular recoil with resultant closure of the artery and/or deposition of blood platelets and fibrin along the damaged length of the newly opened blood vessel.

Restenosis after percutaneous transluminal coronary angioplasty is a more gradual process initiated by vascular injury. Multiple processes, including thrombosis, inflammation, growth factor and cytokine release, cell proliferation, cell migration and extracellular matrix synthesis each contribute to the restenotic process.

While the exact mechanism of restenosis is not completely understood, the general aspects of the restenosis process have been identified. In the normal arterial wall, smooth muscle cells proliferate at a low rate, approximately less than 0.1 percent per day. Smooth muscle cells in the vessel walls exist in a contractile phenotype characterized by eighty to ninety percent of the cell cytoplasmic volume occupied with the contractile apparatus. Endoplasmic reticulum, Golgi, and free ribosomes are few and are located in the perinuclear region. Extracellular matrix surrounds the

2

smooth muscle cells and is rich in heparin-like glycosaminoglycans which are believed to be responsible for maintaining smooth muscle cells in the contractile phenotypic state (Campbell and Campbell, 1985).

Upon pressure expansion of an intracoronary balloon catheter during angioplasty, smooth muscle cells within the vessel wall become injured, initiating a thrombotic and inflammatory response. Cell derived growth factors such as platelet derived growth factor, fibroblast growth factor, epidermal growth factor, thrombin, etc., released from platelets, invading macrophages and/or leukocytes, or directly from the smooth muscle cells provoke proliferative and migratory responses in medial smooth muscle cells. These cells undergo a change from the contractile phenotype to a synthetic phenotype characterized by only a few contractile filament bundles, extensive rough endoplasmic reticulum, Golgi and free ribosomes. Proliferation/migration usually begins within one to two days post-injury and peaks several days thereafter (Campbell and Campbell, 1987; Clowes and Schwartz, 1985).

Daughter cells migrate to the intimal layer of arterial smooth muscle and continue to proliferate and secrete significant amounts of extracellular matrix proteins. Proliferation, migration and extracellular matrix synthesis continue until the damaged endothelial layer is repaired at which time proliferation slows within the intima, usually within seven to fourteen days post-injury. The newly formed tissue is called neointima. The further vascular narrowing that occurs over the next three to six months is due primarily to negative or constrictive remodeling.

Simultaneous with local proliferation and migration, inflammatory cells invade the site of vascular injury. Within three to seven days post-injury, inflammatory cells have migrated to the deeper layers of the vessel wall. In animal models employing either balloon injury or stent implantation, inflammatory cells may persist at the site of vascular injury for at least thirty days (Tanaka et al., 1993; Edelman et al., 1998). Inflammatory cells therefore are present and may contribute to both the acute and chronic phases of restenosis.

Numerous agents have been examined for presumed anti-proliferative actions in restenosis and have shown some activity in experimental animal models. Some of the agents which have been shown to successfully reduce the extent of intimal hyperplasia in animal models include: heparin and heparin fragments (Clowes, A. W. and Karnovsky M., *Nature* 265: 25-26, 1977; Guyton, J. R. et al., *Circ. Res.*, 46: 625-634, 1980; Clowes, A. W. and Clowes, M. M., *Lab. Invest.* 52: 611-616, 1985; Clowes, A. W. and Clowes, M. M., *Circ. Res.* 58: 839-845, 1986; Majesky et al., *Circ. Res.* 61: 296-300, 1987; Snow et al., *Am. J. Pathol.* 137: 313-330, 1990; Okada, T. et al., *Neurosurgery* 25: 92-98, 1989), colchicine (Currier, J. W. et al., *Circ.* 80: 11-66, 1989), taxol (Sollot, S. J. et al., *J. Clin. Invest.* 95: 1869-1876, 1995), angiotensin converting enzyme (ACE) inhibitors (Powell, J. S. et al., *Science*, 245: 186-188, 1989), angiopeptin (Lundergan, C. F. et al. *Am. J. Cardiol.* 17(Suppl. B):132B-136B, 1991), cyclosporin A (Jonasson, L. et al., *Proc. Natl., Acad. Sci.*, 85: 2303, 1988), goat-anti-rabbit PDGF antibody (Ferns, G. A. A., et al., *Science* 253: 1129-1132, 1991), terbinafine (Nemecek, G. M. et al., *J. Pharmacol. Exp. Ther.* 248: 1167-1174, 1989), trapidil (Liu, M. W. et al., *Circ.* 81: 1089-1093, 1990), tranilast (Fukuyama, J. et al., *Eur. J. Pharmacol.* 318: 327-332, 1996), interferon-gamma (Hansson, G. K. and Holm, J., *Circ.* 84: 1266-1272, 1991), rapamycin (Marx, S. O. et al., *Circ. Res.* 76: 412-417, 1995), corticosteroids (Colburn, M. D. et al., *J. Vasc. Surg.* 15:

US 7,300,662 B2

3

510-518, 1992), see also Berk, B. C. et al., *J. Am. Coll. Cardiol.* 17: 111B-117B, 1991), ionizing radiation (Weinberger, J. et al., *Int. J. Rad. Onc. Biol. Phys.* 36: 767-775, 1996), fusion toxins (Farb, A. et al., *Circ. Res.* 80: 542-550, 1997) antisense oligonucleotides (Simons, M. et al., *Nature* 359: 67-70, 1992) and gene vectors (Chang, M. W. et al., *J. Clin. Invest.* 96: 2260-2268, 1995). Anti-proliferative effects on smooth muscle cells in vitro have been demonstrated for many of these agents, including heparin and heparin conjugates, taxol, tranilast, colchicine, ACE inhibitors, fusion toxins, antisense oligonucleotides, rapamycin and ionizing radiation. Thus, agents with diverse mechanisms of smooth muscle cell inhibition may have therapeutic utility in reducing intimal hyperplasia.

However, in contrast to animal models, attempts in human angioplasty patients to prevent restenosis by systemic pharmacologic means have thus far been unsuccessful. Neither aspirin-dipyridamole, ticlopidine, anti-coagulant therapy (acute heparin, chronic warfarin, hirudin or hirulog), thromboxane receptor antagonism nor steroids have been effective in preventing restenosis, although platelet inhibitors have been effective in preventing acute reocclusion after angioplasty (Mak and Topol, 1997; Lang et al., 1991; Popma et al., 1991). The platelet GP IIb/IIIa receptor, antagonist, Reopro is still under study but has not shown promising results for the reduction in restenosis following angioplasty and stenting. Other agents, which have also been unsuccessful in the prevention of restenosis, include the calcium channel antagonists, prostacyclin mimetics, angiotensin converting enzyme inhibitors, serotonin receptor antagonists, and anti-proliferative agents. These agents must be given systemically, however, and attainment of a therapeutically effective dose may not be possible; anti-proliferative (or anti-restenosis) concentrations may exceed the known toxic concentrations of these agents so that levels sufficient to produce smooth muscle inhibition may not be reached (Mak and Topol, 1997; Lang et al., 1991; Popma et al., 1991).

Additional clinical trials in which the effectiveness for preventing restenosis utilizing dietary fish oil supplements or cholesterol lowering agents has been examined showing either conflicting or negative results so that no pharmacological agents are as yet clinically available to prevent post-angioplasty restenosis (Mak and Topol, 1997; Franklin and Faxon, 1993; Serruys, P. W. et al., 1993). Recent observations suggest that the antilipid/antioxidant agent, probucol may be useful in preventing restenosis but this work requires confirmation (Tardif et al., 1997; Yokoi, et al., 1997). Probuco is presently not approved for use in the United States and a thirty-day pretreatment period would preclude its use in emergency angioplasty. Additionally, the application of ionizing radiation has shown significant promise in reducing or preventing restenosis after angioplasty in patients with stents (Teirstein et al., 1997). Currently, however, the most effective treatments for restenosis are repeat angioplasty, atherectomy or coronary artery bypass grafting, because no therapeutic agents currently have Food and Drug Administration approval for use for the prevention of post-angioplasty restenosis.

Unlike systemic pharmacologic therapy, stents have proven effective in significantly reducing restenosis. Typically, stents are balloon-expandable slotted metal tubes (usually, but not limited to, stainless steel), which, when expanded within the lumen of an angioplastied coronary artery, provide structural support through rigid scaffolding to the arterial wall. This support is helpful in maintaining vessel lumen patency. In two randomized clinical trials,

4

stents increased angiographic success after percutaneous transluminal coronary angioplasty, by increasing minimal lumen diameter and reducing, but not eliminating, the incidence of restenosis at six months (Serruys et al., 1994; Fischman et al., 1994).

Additionally, the heparin coating of stents appears to have the added benefit of producing a reduction in sub-acute thrombosis after stent implantation (Serruys et al., 1996). Thus, sustained mechanical expansion of a stenosed coronary artery with a stent has been shown to provide some measure of restenosis prevention, and the coating of stents with heparin has demonstrated both the feasibility and the clinical usefulness of delivering drugs locally, at the site of injured tissue.

Accordingly, there exists a need for effective drugs and drug delivery systems for the effective prevention and treatment of neointimal thickening that occurs after percutaneous transluminal coronary angioplasty and stent implantation.

#### SUMMARY OF THE INVENTION

The drugs and drug delivery systems of the present invention provide a means for overcoming the difficulties associated with the methods and devices currently in use as briefly described above.

In accordance with one aspect, the present invention is directed to a method for the prevention of constrictive remodeling. The method comprises the controlled delivery, by release from an intraluminal medical device, of a compound in therapeutic dosage amounts.

In accordance with another aspect, the present invention is directed to a drug delivery device. The drug delivery device comprises an intraluminal medical device and a therapeutic dosage of an agent releasably affixed to the intraluminal medical device for the treatment of constrictive vascular remodeling.

The drugs and drug delivery systems of the present invention utilize a stent or graft in combination with rapamycin or other drugs/agents/compounds to prevent and treat neointimal hyperplasia, i.e. restenosis, following percutaneous transluminal coronary angioplasty and stent implantation. It has been determined that rapamycin functions to inhibit smooth muscle cell proliferation through a number of mechanisms. It has also been determined that rapamycin eluting stent coatings produce superior effects in humans, when compared to animals, with respect to the magnitude and duration of the reduction in neointimal hyperplasia. Rapamycin administration from a local delivery platform also produces an anti-inflammatory effect in the vessel wall that is distinct from and complimentary to its smooth muscle cell anti-proliferative effect. In addition, it has also been demonstrated that rapamycin inhibits constrictive vascular remodeling in humans.

Other drugs, agents or compounds which mimic certain actions of rapamycin may also be utilized in combination with local delivery systems or platforms.

The local administration of drugs, agents or compounds to stented vessels have the additional therapeutic benefit of higher tissue concentration than that which would be achievable through the systemic administration of the same drugs, agents or compounds. Other benefits include reduced systemic toxicity, single treatment, and ease of administration. An additional benefit of a local delivery device and drug, agent or compound therapy may be to reduce the dose of the therapeutic drugs, agents or compounds and thus limit their toxicity, while still achieving a reduction in restenosis.

US 7,300,662 B2

5

## BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other features and advantages of the invention will be apparent from the following, more particular description of preferred embodiments of the invention, as illustrated in the accompanying drawings.

FIG. 1 is a chart indicating the effectiveness of rapamycin as an anti-inflammatory relative to other anti-inflammatories.

FIG. 2 is a view along the length of a stent (ends not shown) prior to expansion showing the exterior surface of the stent and the characteristic banding pattern.

FIG. 3 is a perspective view of the stent of FIG. 1 having reservoirs in accordance with the present invention.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

As stated above, the proliferation of vascular smooth muscle cells in response to mitogenic stimuli that are released during balloon angioplasty and stent implantation is the primary cause of neointimal hyperplasia. Excessive neointimal hyperplasia can often lead to impairment of blood flow, cardiac ischemia and the need for a repeat intervention in selected patients in high risk treatment groups. Yet repeat revascularization incurs risk of patient morbidity and mortality while adding significantly to the cost of health care. Given the widespread use of stents in interventional practice, there is a clear need for safe and effective inhibitors of neointimal hyperplasia.

Rapamycin is a macrocyclic triene antibiotic produced by streptomyces hygroscopicus as disclosed in U.S. Pat. No. 3,929,992. It has been found that rapamycin inhibits the proliferation of vascular smooth muscle cells in vivo. Accordingly, rapamycin may be utilized in treating intimal smooth muscle cell hyperplasia, restenosis and vascular occlusion in a mammal, particularly following either biologically or mechanically mediated vascular injury, or under conditions that would predispose a mammal to suffering such a vascular injury. Rapamycin functions to inhibit smooth muscle cell proliferation and does not interfere with the re-endothelialization of the vessel walls.

Rapamycin functions to inhibit smooth muscle cell proliferation through a number of mechanisms. In addition, rapamycin reduces the other effects caused by vascular injury, for example, inflammation. The operation and various functions of rapamycin are described in detail below. Rapamycin as used throughout this application shall include rapamycin, rapamycin analogs, derivatives and congeners that bind FKBP12 and possess the same pharmacologic properties as rapamycin.

Rapamycin reduces vascular hyperplasia by antagonizing smooth muscle proliferation in response to mitogenic signals that are released during angioplasty. Inhibition of growth factor and cytokine mediated smooth muscle proliferation at the late G1 phase of the cell cycle is believed to be the dominant mechanism of action of rapamycin. However, rapamycin is also known to prevent T-cell proliferation and differentiation when administered systemically. This is the basis for its immunosuppressive activity and its ability to prevent graft rejection.

The molecular events that are responsible for the actions of rapamycin, a known anti-proliferative, which acts to reduce the magnitude and duration of neointimal hyperplasia, are still being elucidated. It is known, however, that rapamycin enters cells and binds to a high-affinity cytosolic protein called FKBP12. The complex of rapamycin and

6

FKBP12 in turn binds to and inhibits a phosphoinositide (PI)-3 kinase called the "mammalian Target of Rapamycin" or TOR. TOR is a protein kinase that plays a key role in mediating the downstream signaling events associated with mitogenic growth factors and cytokines in smooth muscle cells and T lymphocytes. These events include phosphorylation of p27, phosphorylation of p70 s6 kinase and phosphorylation of 4BP-1, an important regulator of protein translation.

It is recognized that rapamycin reduces restenosis by inhibiting neointimal hyperplasia. However, there is evidence that rapamycin may also inhibit the other major component of restenosis, namely, negative remodeling. Remodeling is a process whose mechanism is not clearly understood but which results in shrinkage of the external elastic lamina and reduction in luminal area over time, generally a period of approximately three to six months in humans.

Negative or constrictive vascular remodeling may be quantified angiographically as the percent diameter stenosis at the lesion site where there is no stent to obstruct the process. If late lumen loss is abolished in-lesion, it may be inferred that negative remodeling has been inhibited. Another method of determining the degree of remodeling involves measuring in-lesion external elastic lamina area using intravascular ultrasound (IVUS). Intravascular ultrasound is a technique that can image the external elastic lamina as well as the vascular lumen. Changes in the external elastic lamina proximal and distal to the stent from the post-procedural timepoint to four-month and twelve-month follow-ups are reflective of remodeling changes.

Evidence that rapamycin exerts an effect on remodeling comes from human implant studies with rapamycin coated stents showing a very low degree of restenosis in-lesion as well as in-stent. In-lesion parameters are usually measured approximately five millimeters on either side of the stent i.e. proximal and distal. Since the stent is not present to control remodeling in these zones which are still affected by balloon expansion, it may be inferred that rapamycin is preventing vascular remodeling.

The data in Table 1 below illustrate that in-lesion percent diameter stenosis remains low in the rapamycin treated groups, even at twelve months. Accordingly, these results support the hypothesis that rapamycin reduces remodeling.

TABLE 1.0

Angiographic In-Lesion Percent Diameter Stenosis (%, mean $\pm$ SD and "n=") In Patients Who Received a Rapamycin-Coated Stent			
Coating Group	Post Placement	4-6 month Follow Up	12 month Follow Up
Brazil	10.6 $\pm$ 5.7 (30)	13.6 $\pm$ 8.6 (30)	22.3 $\pm$ 7.2 (15)
Netherlands	14.7 $\pm$ 8.8	22.4 $\pm$ 6.4	—

Additional evidence supporting a reduction in negative remodeling with rapamycin comes from intravascular ultrasound data that was obtained from a first-in-man clinical program as illustrated in Table 2 below.

US 7,300,662 B2

7

TABLE 2.0

Matched IVUS data in Patients Who Received a Rapamycin-Coated Stent			
IVUS Parameter	Post (n=)	4-Month Follow-Up (n=)	12-Month Follow-Up (n=)
Mean proximal vessel area (mm <sup>2</sup> )	16.53 ± 3.53 (27)	16.31 ± 4.36 (28)	13.96 ± 2.26 (13)
Mean distal vessel area (mm <sup>2</sup> )	13.12 ± 3.68 (26)	13.53 ± 4.17 (26)	12.49 ± 3.25 (14)

The data illustrated that there is minimal loss of vessel area proximally or distally which indicates that inhibition of negative remodeling has occurred in vessels treated with rapamycin-coated stents.

Other than the stent itself, there have been no effective solutions to the problem of vascular remodeling. Accordingly, rapamycin may represent a biological approach to controlling the vascular remodeling phenomenon.

It may be hypothesized that rapamycin acts to reduce negative remodeling in several ways. By specifically blocking the proliferation of fibroblasts in the vascular wall in response to injury, rapamycin may reduce the formation of vascular scar tissue. Rapamycin may also affect the translation of key proteins involved in collagen formation or metabolism.

Rapamycin used in this context includes rapamycin and all analogs, derivatives and congeners that bind FKBP12 and possess the same pharmacologic properties as rapamycin.

8

In a preferred embodiment, the rapamycin is delivered by a local delivery device to control negative remodeling of an arterial segment after balloon angioplasty as a means of reducing or preventing restenosis. While any delivery device may be utilized, it is preferred that the delivery device comprises a stent that includes a coating or sheath which elutes or releases rapamycin. The delivery system for such a device may comprise a local infusion catheter that delivers rapamycin at a rate controlled by the administrator.

Rapamycin may also be delivered systemically using an oral dosage form or a chronic injectable depot form or a patch to deliver rapamycin for a period ranging from about seven to forty-five days to achieve vascular tissue levels that are sufficient to inhibit negative remodeling. Such treatment is to be used to reduce or prevent restenosis when administered several days prior to elective angioplasty with or without a stent.

Data generated in porcine and rabbit models show that the release of rapamycin into the vascular wall from a nonerodible polymeric stent coating in a range of doses (35-430 ug/5-18 mm coronary stent) produces a peak fifty to fifty-five percent reduction in neointimal hyperplasia as set forth in Table 3 below. This reduction, which is maximal at about twenty-eight to thirty days, is typically not sustained in the range of ninety to one hundred eighty days in the porcine model as set forth in Table 4 below.

TABLE 3.0

Animal Studies with Rapamycin-coated stents.						
Values are mean ± Standard Error of Mean						
Study	Duration	Stent <sup>1</sup>	Neointimal Area		% Change From	
			Rapamycin	N (mm <sup>2</sup> )	Polyme	Metal
Porcine						
98009	14 days	Metal		8 2.04 ± 0.17		
		1X + rapamycin	153 µg	8 1.66 ± 0.17*	-42%	-19%
		1X + TC300 + rapamycin	155 µg	8 1.51 ± 0.19*	-47%	-26%
99005	28 days	Metal		10 2.29 ± 0.21		
				9 3.91 ± 0.60**		
		1X + TC30 + rapamycin	130 µg	8 2.81 ± 0.34		+23%
99006	28 days	1X + TC100 + rapamycin	120 µg	9 2.62 ± 0.21		+14%
		Metal		12 4.57 ± 0.46		
		EVA/BMA 3X		12 5.02 ± 0.62		+10%
		1X + rapamycin	125 µg	11 2.84 ± 0.31* **	-43%	-38%
		3X + rapamycin	430 µg	12 3.06 ± 0.17* **	-39%	-33%
		3X + rapamycin	157 µg	12 2.77 ± 0.41* **	-45%	-39%
99011	28 days	Metal		11 3.09 ± 0.27		
				11 4.52 ± 0.37		
		1X + rapamycin	189 µg	14 3.05 ± 0.35		-1%
99021	60 days	3X + rapamycin/dex	182/363 µg	14 2.72 ± 0.71		-12%
		Metal		12 2.14 ± 0.25		
		1X + rapamycin	181 µg	12 2.95 ± 0.38		+38%
99034	28 days	Metal		8 5.24 ± 0.58		
		1X + rapamycin	186 µg	8 2.47 ± 0.33**		-53%
		3X + rapamycin/dex	185/369 µg	6 2.42 ± 0.64**		-54%
20001	28 days	Metal		6 1.81 ± 0.09		
20007	30 days	1X + rapamycin	172 µg	5 1.66 ± 0.44		-8%
		Metal		9 2.94 ± 0.43		
		1XTC + rapamycin	155 µg	10 1.40 ± 0.11*		-52%
Rabbit						
99019	28 days	Metal		8 1.20 ± 0.07		

A1251



US 7,300,662 B2

9

10

TABLE 3.0-continued

Animal Studies with Rapamycin-coated stents. Values are mean $\pm$ Standard Error of Mean					
Study	Duration	Stent <sup>1</sup>	Rapamycin	Neointimal Area	% Change From
				N (mm <sup>2</sup> )	Polyme Metal
99020	28 days	EVA/BMA 1X		10 1.26 $\pm$ 0.16	+5%
		1X + rapamycin	64 $\mu$ g	9 0.92 $\pm$ 0.14	-27%
		1X + rapamycin	196 $\mu$ g	10 0.66 $\pm$ 0.12* **	-48%
		Metal		12 1.18 $\pm$ 0.10	-45%
		EVA/BMA 1X + rapamycin	197 $\mu$ g	8 0.81 $\pm$ 0.16	-32%

<sup>1</sup>Stent nomenclature: EVA/BMA 1X, 2X, and 3X signifies approx. 500  $\mu$ g, 1000  $\mu$ g, and 1500  $\mu$ g total mass (polymer + drug), respectively. TC, top coat of 30  $\mu$ g, 100  $\mu$ g, or 300  $\mu$ g drug-free BMA; Biphasic; 2  $\times$  1X layers of rapamycin in EVA/BMA separated by a 100  $\mu$ g drug-free BMA layer.

<sup>2</sup>0.25 mg/kg/d  $\times$  14 d preceded by a loading dose of 0.5 mg/kg/d  $\times$  3 d prior to stent implantation.

\*p < 0.05 from EVA/BMA control.

\*\*p < 0.05 from Metal;

<sup>#</sup>Inflammation score: (0 = essentially no intimal involvement; 1 = <25% intima involved; 2 =  $\geq$ 25% intima involved; 3 = >50% intima involved).

TABLE 4.0

180 day Porcine Study with Rapamycin-coated stents.								
Values are mean ± Standard Error of Mean								
Study	Duration	Stent <sup>1</sup>	Rapamycin	N	Neointimal Area	% Change From	Inflammation	
					(mm <sup>2</sup> )	Polyme Metal	Score #	
20007 (ETP-2-002233-P)	3 days	Metal		10	0.38 ± 0.06			1.05 ± 0.06
		1XTC + rapamycin	155 µg	10	0.29 ± 0.03	-24%		1.08 ± 0.04
	30 days	Metal		9	2.94 ± 0.43			0.11 ± 0.08
		1XTC + rapamycin	155 µg	10	1.40 ± 0.11*	-52%*		0.25 ± 0.10
	90 days	Metal		10	3.45 ± 0.34			0.20 ± 0.08
1XTC + rapamycin		155 µg	10	3.03 ± 0.29	-12%		0.80 ± 0.23	
1X + rapamycin		171 µg	10	2.86 ± 0.35	-17%		0.60 ± 0.23	
180 days	Metal		10	3.65 ± 0.39			0.65 ± 0.21	
	1XTC + rapamycin		155 µg	10	3.34 ± 0.31	-8%		1.50 ± 0.34
	1X + rapamycin		171 µg	10	3.87 ± 0.28	+6%		1.68 ± 0.37

The release of rapamycin into the vascular wall of a human from a nonerodible polymeric stent coating provides superior results with respect to the magnitude and duration of the reduction in neointimal hyperplasia within the stent as compared to the vascular walls of animals as set forth above.

Humans implanted with a rapamycin coated stent comprising rapamycin in the same dose range as studied in animal models using the same polymeric matrix, as

described above, reveal a much more profound reduction in neointimal hyperplasia than observed in animal models, based on the magnitude and duration of reduction in neointima. The human clinical response to rapamycin reveals essentially total abolition of neointimal hyperplasia inside the stent using both angiographic and intravascular ultrasound measurements. These results are sustained for at least one year as set forth in Table 5 below.

TABLE 5.0

Patients Treated (N = 45 patients) with a Rapamycin-coated Stent		
Effectiveness Measures	Sirolimus FIM (N = 45 Patients, 45 Lesions)	95% Confidence Limit
Procedure Success (QCA)	100.0% (45/45)	[92.1%, 100.0%]
4-month In-Stent Diameter Stenosis (%)		
Mean $\pm$ SD (N)	4.8% $\pm$ 6.1% (30)	[2.6%, 7.0%]
Range (min, max)	(-8.2%, 14.9%)	
6-month In-Stent Diameter Stenosis (%)		
Mean $\pm$ SD (N)	8.9% $\pm$ 7.6% (13)	[4.8%, 13.0%]
Range (min, max)	(-2.9%, 20.4%)	
12-month In-Stent Diameter Stenosis (%)		
Mean $\pm$ SD (N)	8.9% $\pm$ 6.1% (15)	[5.8%, 12.0%]
Range (min, max)	(-3.0%, 22.0%)	

A1252

US 7,300,662 B2

11

12

TABLE 5.0-continued

Patients Treated (N = 45 patients) with a Rapamycin-coated Stent		
Effectiveness Measures	Sirolimus FIM (N = 45 Patients, 45 Lesions)	95% Confidence Limit
<u>4-month In-Stent Late Loss (mm)</u>		
Mean $\pm$ SD (N)	0.00 $\pm$ 0.29 (30)	[-0.10, 0.10]
Range (min, max)	(-0.51, 0.45)	
<u>6-month In-Stent Late Loss (mm)</u>		
Mean $\pm$ SD (N)	0.25 $\pm$ 0.27 (13)	[0.10, 0.39]
Range (min, max)	(-0.51, 0.91)	
<u>12-month In-Stent Late Loss (mm)</u>		
Mean $\pm$ SD (N)	0.11 $\pm$ 0.36 (15)	[-0.08, 0.29]
Range (min, max)	(-0.51, 0.82)	
<u>4-month Obstruction Volume (%) (IVUS)</u>		
Mean $\pm$ SD (N)	10.48% $\pm$ 2.78% (28)	[9.45%, 11.51%]
Range (min, max)	(4.60%, 16.35%)	
<u>6-month Obstruction Volume (%) (IVUS)</u>		
Mean $\pm$ SD (N)	7.22% $\pm$ 4.60% (13)	[4.72%, 9.72%]
Range (min, max)	(3.82%, 19.88%)	
<u>12-month Obstruction Volume (%) (IVUS)</u>		
Mean $\pm$ SD (N)	2.11% $\pm$ 5.28% (15)	[0.00%, 4.78%]
Range (min, max)	(0.00%, 19.89%)	
6-month Target Lesion Revascularization (TLR)	0.0% (0/30)	[0.0%, 9.5%]
12-month Target Lesion Revascularization (TLR)	0.0% (0/15)	[0.0%, 18.1%]

QCA = Quantitative Coronary Angiography

SD = Standard Deviation

IVUS = Intravascular Ultrasound

Rapamycin produces an unexpected benefit in humans when delivered from a stent by causing a profound reduction in in-stent neointimal hyperplasia that is sustained for at least one year. The magnitude and duration of this benefit in humans is not predicted from animal model data. Rapamycin used in this context includes rapamycin and all analogs, derivatives and congeners that bind FKBP12 and possess the same pharmacologic properties as rapamycin.

These results may be due to a number of factors. For example, the greater effectiveness of rapamycin in humans is due to greater sensitivity of its mechanism(s) of action toward the pathophysiology of human vascular lesions compared to the pathophysiology of animal models of angioplasty. In addition, the combination of the dose applied to the stent and the polymer coating that controls the release of the drug is important in the effectiveness of the drug.

As stated above, rapamycin reduces vascular hyperplasia by antagonizing smooth muscle proliferation in response to mitogenic signals that are released during angioplasty injury. Also, it is known that rapamycin prevents T-cell proliferation and differentiation when administered systemically. It has also been determined that rapamycin exerts a local inflammatory effect in the vessel wall when administered from a stent in low doses for a sustained period of time (approximately two to six weeks). The local anti-inflammatory benefit is profound and unexpected. In combination with the smooth muscle anti-proliferative effect, this dual mode of action of rapamycin may be responsible for its exceptional efficacy.

Accordingly, rapamycin delivered from a local device platform, reduces neointimal hyperplasia by a combination of anti-inflammatory and smooth muscle anti-proliferative effects. Rapamycin used in this context means rapamycin

and all analogs, derivatives and congeners that bind FKBP12 and possess the same pharmacologic properties as rapamycin. Local device platforms include stent coatings, stent sheaths, grafts and local drug infusion catheters or porous balloons or any other suitable means for the in situ or local delivery of drugs, agents or compounds.

The anti-inflammatory effect of rapamycin is evident in data from an experiment, illustrated in Table 6, in which rapamycin delivered from a stent was compared with dexamethasone delivered from a stent. Dexamethasone, a potent steroidal anti-inflammatory agent, was used as a reference standard. Although dexamethasone is able to reduce inflammation scores, rapamycin is far more effective than dexamethasone in reducing inflammation scores. In addition, rapamycin significantly reduces neointimal hyperplasia, unlike dexamethasone.

TABLE 6.0

Group				
Rapamycin	N=	Neointimal Area (mm <sup>2</sup> )	% Area Stenosis	Inflammation Score
Rap				
Uncoated	8	5.24 $\pm$ 1.65	54 $\pm$ 19	0.97 $\pm$ 1.00
Dexamethasone (Dex)	8	4.31 $\pm$ 3.02	45 $\pm$ 31	0.39 $\pm$ 0.24
Rapamycin (Rap)	7	2.47 $\pm$ 0.94*	26 $\pm$ 10*	0.13 $\pm$ 0.19*
Rap + Dex	6	2.42 $\pm$ 1.58*	26 $\pm$ 18*	0.17 $\pm$ 0.30*

\* = significance level P &lt; 0.05

Rapamycin has also been found to reduce cytokine levels in vascular tissue when delivered from a stent. The data in FIG. 1 illustrates that rapamycin is highly effective in reducing monocyte chemotactic protein (MCP-1) levels in

A1253

US 7,300,662 B2

13

the vascular wall. MCP-1 is an example of a proinflammatory/chemotactic cytokine that is elaborated during vessel injury. Reduction in MCP-1 illustrates the beneficial effect of rapamycin in reducing the expression of proinflammatory mediators and contributing to the anti-inflammatory effect of rapamycin delivered locally from a stent. It is recognized that vascular inflammation in response to injury is a major contributor to the development of neointimal hyperplasia.

Since rapamycin may be shown to inhibit local inflammatory events in the vessel it is believed that this could explain the unexpected superiority of rapamycin in inhibiting neointima.

As set forth above, rapamycin functions on a number of levels to produce such desired effects as the prevention of T-cell proliferation, the inhibition of negative remodeling, the reduction of inflammation, and the prevention of smooth muscle cell proliferation. While the exact mechanisms of these functions are not completely known, the mechanisms that have been identified may be expanded upon.

Studies with rapamycin suggest that the prevention of smooth muscle cell proliferation by blockade of the cell cycle is a valid strategy for reducing neointimal hyperplasia. Dramatic and sustained reductions in late lumen loss and neointimal plaque volume have been observed in patients receiving rapamycin delivered locally from a stent. The present invention expands upon the mechanism of rapamycin to include additional approaches to inhibit the cell cycle and reduce neointimal hyperplasia without producing toxicity.

The cell cycle is a tightly controlled biochemical cascade of events that regulate the process of cell replication. When cells are stimulated by appropriate growth factors, they move from G<sub>0</sub> (quiescence) to the G<sub>1</sub> phase of the cell cycle. Selective inhibition of the cell cycle in the G<sub>1</sub> phase, prior to DNA replication (S phase), may offer therapeutic advantages of cell preservation and viability while retaining anti-proliferative efficacy when compared to therapeutics that act later in the cell cycle i.e. at S, G<sub>2</sub> or M phase.

Accordingly, the prevention of intimal hyperplasia in blood vessels and other conduit vessels in the body may be achieved using cell cycle inhibitors that act selectively at the G<sub>1</sub> phase of the cell cycle. These inhibitors of the G<sub>1</sub> phase of the cell cycle may be small molecules, peptides, proteins, oligonucleotides or DNA sequences. More specifically, these drugs or agents include inhibitors of cyclin dependent kinases (cdk's) involved with the progression of the cell cycle through the G<sub>1</sub> phase, in particular cdk2 and cdk4.

Examples of drugs, agents or compounds that act selectively at the G<sub>1</sub> phase of the cell cycle include small molecules such as flavopiridol and its structural analogs that have been found to inhibit cell cycle in the late G<sub>1</sub> phase by antagonism of cyclin dependent kinases. Therapeutic agents that elevate an endogenous kinase inhibitory protein<sup>kip</sup> called P27, sometimes referred to as P27<sup>kip1</sup>, that selectively inhibits cyclin dependent kinases may be utilized. This includes small molecules, peptides and proteins that either block the degradation of P27 or enhance the cellular production of P27, including gene vectors that can transfect the gene to produce P27. Staurosporin and related small molecules that block the cell cycle by inhibiting protein kinases may be utilized. Protein kinase inhibitors, including the class of tyrphostins that selectively inhibit protein kinases to antagonize signal transduction in smooth muscle in response to a broad range of growth factors such as PDGF and FGF may also be utilized.

Any of the drugs, agents or compounds discussed above may be administered either systemically, for example,

14

orally, intravenously, intramuscularly, subcutaneously, nasally or intradermally, or locally, for example, stent coating, stent covering or local delivery catheter. In addition, the drugs or agents discussed above may be formulated for fast-release or slow release with the objective of maintaining the drugs or agents in contact with target tissues for a period ranging from three days to eight weeks.

As set forth above, the complex of rapamycin and FKBP12 binds to and inhibits a phosphoinositide (PI)-3 kinase called the mammalian Target of Rapamycin or TOR. An antagonist of the catalytic activity of TOR, functioning as either an active site inhibitor or as an allosteric modulator, i.e. an indirect inhibitor that allosterically modulates, would mimic the actions of rapamycin but bypass the requirement for FKBP12. The potential advantages of a direct inhibitor of TOR include better tissue penetration and better physical/chemical stability. In addition, other potential advantages include greater selectivity and specificity of action due to the specificity of an antagonist for one of multiple isoforms of TOR that may exist in different tissues, and a potentially different spectrum of downstream effects leading to greater drug efficacy and/or safety.

The inhibitor may be a small organic molecule (approximate mw<1000), which is either a synthetic or naturally derived product. Wortmanin may be an agent which inhibits the function of this class of proteins. It may also be a peptide or an oligonucleotide sequence. The inhibitor may be administered either systemically (orally, intravenously, intramuscularly, subcutaneously, nasally, or intradermally) or locally (stent coating, stent covering, local drug delivery catheter). For example, the inhibitor may be released into the vascular wall of a human from a nonerodible polymeric stent coating. In addition, the inhibitor may be formulated for fast-release or slow release with the objective of maintaining the rapamycin or other drug, agent or compound in contact with target tissues for a period ranging from three days to eight weeks.

As stated previously, the implantation of a coronary stent in conjunction with balloon angioplasty is highly effective in treating acute vessel closure and may reduce the risk of restenosis. Intravascular ultrasound studies (Mintz et al., 1996) suggest that coronary stenting effectively prevents vessel constriction and that most of the late luminal loss after stent implantation is due to plaque growth, probably related to neointimal hyperplasia. The late luminal loss after coronary stenting is almost two times higher than that observed after conventional balloon angioplasty. Thus, inasmuch as stents prevent at least a portion of the restenosis process, the use of drugs, agents or compounds which prevent inflammation and proliferation, or prevent proliferation by multiple mechanisms, combined with a stent may provide the most efficacious treatment for post-angioplasty restenosis.

The local delivery of drugs, agents or compounds from a stent has the following advantages; namely, the prevention of vessel recoil and remodeling through the scaffolding action of the stent and the drugs, agents or compounds and the prevention of multiple components of neointimal hyperplasia. This local administration of drugs, agents or compounds to stented coronary arteries may also have additional therapeutic benefit. For example, higher tissue concentrations would be achievable than that which would occur with systemic administration, reduced systemic toxicity, and single treatment and ease of administration. An additional benefit of drug therapy may be to reduce the dose of the therapeutic compounds, thereby limiting their toxicity, while still achieving a reduction in restenosis.

A1254

US 7,300,662 B2

15

There are a multiplicity of different stents that may be utilized following percutaneous transluminal coronary angioplasty. Although any number of stents may be utilized in accordance with the present invention, for simplicity, one particular stent will be described in exemplary embodiments of the present invention. The skilled artisan will recognize that any number of stents may be utilized in connection with the present invention.

A stent is commonly used as a tubular structure left inside the lumen of a duct to relieve an obstruction. Commonly, stents are inserted into the lumen in a non-expanded form and are then expanded autonomously, or with the aid of a second device in situ. A typical method of expansion occurs through the use of a catheter-mounted angioplasty balloon which is inflated within the stenosed vessel or body passageway in order to shear and disrupt the obstructions associated with the wall components of the vessel and to obtain an enlarged lumen. As set forth below, self-expanding stents may also be utilized.

FIG. 2 illustrates an exemplary stent 100 which may be utilized in accordance with an exemplary embodiment of the present invention. The expandable cylindrical stent 100 comprises a fenestrated structure for placement in a blood vessel, duct or lumen to hold the vessel, duct or lumen open, more particularly for protecting a segment of artery from restenosis after angioplasty. The stent 100 may be expanded circumferentially and maintained in an expanded configuration, that is circumferentially or radially rigid. The stent 100 is axially flexible and when flexed at a band, the stent 100 avoids any externally-protruding component parts.

The stent 100 generally comprises first and second ends with an intermediate section therebetween. The stent 100 has a longitudinal axis and comprises a plurality of longitudinally disposed bands 102, wherein each band 102 defines a generally continuous wave along a line segment parallel to the longitudinal axis. A plurality of circumferentially arranged links 104 maintain the bands 102 in a substantially tubular structure. Essentially, each longitudinally disposed band 102 is connected at a plurality of periodic locations, by a short circumferentially arranged link 104 to an adjacent band 102. The wave associated with each of the bands 102 has approximately the same fundamental spatial frequency in the intermediate section, and the bands 102 are so disposed that the wave associated with them are generally aligned so as to be generally in phase with one another. As illustrated in the figure, each longitudinally arranged band 102 undulates through approximately two cycles before there is a link to an adjacent band.

The stent 100 may be fabricated utilizing any number of methods. For example, the stent 100 may be fabricated from a hollow or formed stainless steel tube that may be machined using lasers, electric discharge milling, chemical etching or other means. The stent 100 is inserted into the body and placed at the desired site in an unexpanded form. In one embodiment, expansion may be effected in a blood vessel by a balloon catheter, where the final diameter of the stent 100 is a function of the diameter of the balloon catheter used.

It should be appreciated that a stent 100 in accordance with the present invention may be embodied in a shape-memory material, including, for example, an appropriate alloy of nickel and titanium. In this embodiment, after the stent 100 has been formed it may be compressed so as to occupy a space sufficiently small as to permit its insertion in a blood vessel or other tissue by insertion means, wherein the insertion means include a suitable catheter, or flexible rod. On emerging from the catheter, the stent 100 may be configured to expand into the desired configuration where

16

the expansion is automatic or triggered by a change in pressure, temperature or electrical stimulation.

FIG. 3 illustrates an exemplary embodiment of the present invention utilizing the stent 100 illustrated in FIG. 2. As illustrated, the stent 100 may be modified to comprise a reservoir 106. Each of the reservoirs may be opened or closed as desired. These reservoirs 106 may be specifically designed to hold the drug, agent, compound or combinations thereof to be delivered. Regardless of the design of the stent 100, it is preferable to have the drug, agent, compound or combinations thereof dosage applied with enough specificity and a sufficient concentration to provide an effective dosage in the lesion area. In this regard, the reservoir size in the bands 102 is preferably sized to adequately apply the drug/drug combination dosage at the desired location and in the desired amount.

In an alternate exemplary embodiment, the entire inner and outer surface of the stent 100 may be coated with various drug and drug combinations in therapeutic dosage amounts. A detailed description of exemplary coating techniques is described below.

Rapamycin or any of the drugs, agents or compounds described above may be incorporated into or affixed to the stent in a number of ways and utilizing any number of biocompatible materials. In the exemplary embodiment, the rapamycin is directly incorporated into a polymeric matrix and sprayed onto the outer surface of the stent. The rapamycin elutes from the polymeric matrix over time and enters the surrounding tissue. The rapamycin preferably remains on the stent for at least three days up to approximately six months and more preferably between seven and thirty days.

Any number of non-erodible polymers may be utilized in conjunction with rapamycin. In the exemplary embodiment, the polymeric matrix comprises two layers. The base layer comprises a solution of ethylene-co-vinylacetate and polybutylmethacrylate. The rapamycin is incorporated into this layer. The outer layer comprises only polybutylmethacrylate and acts as a diffusion barrier to prevent the rapamycin from eluting too quickly and entering the surrounding tissues. The thickness of the outer layer or top coat determines the rate at which the rapamycin elutes from the matrix. Essentially, the rapamycin elutes from the matrix by diffusion through the polymer molecules. Polymers tend to move, thereby allowing solids, liquids and gases to escape therefrom. The total thickness of the polymeric matrix is in the range from about 1 micron to about 20 microns or greater. In a preferred exemplary embodiment, the base layer, including the polymer and drug, has a thickness in the range from about 8 microns to about 12 microns and the outer layer has a thickness in the range from about 1 micron to about 2 microns.

The ethylene-co-vinylacetate, polybutylmethacrylate and rapamycin solution may be incorporated into or onto the stent in a number of ways. For example, the solution may be sprayed onto the stent or the stent may be dipped into the solution. In a preferred embodiment, the solution is sprayed onto the stent and then allowed to dry. In another exemplary embodiment, the solution may be electrically charged to one polarity and the stent electrically changed to the opposite polarity. In this manner, the solution and stent will be attracted to one another. In using this type of spraying process, waste may be reduced and more control over the thickness of the coat may be achieved.

Since rapamycin works by entering the surrounding tissue, it is preferably only affixed to the surface of the stent making contact with one tissue. Typically, only the outer surface of the stent makes contact with the tissue. Accord-

A1255

US 7,300,662 B2

17

ingly, in a preferred embodiment, only the outer surface of the stent is coated with rapamycin. For other drugs, agents or compounds, the entire stent may be coated.

It is important to note that different polymers may be utilized for different stents. For example, in the above-described embodiment, ethylene-co-vinylacetate and polybutylmethacrylate are utilized to form the polymeric matrix. This matrix works well with stainless steel stents. Other polymers may be utilized more effectively with stents formed from other materials, including materials that exhibit superelastic properties such as alloys of nickel and titanium.

Although shown and described is what is believed to be the most practical and preferred embodiments, it is apparent that departures from specific designs and methods described and shown will suggest themselves to those skilled in the art and may be used without departing from the spirit and scope of the invention. The present invention is not restricted to the particular constructions described and illustrated, but should be constructed to cohere with all modifications that may fall within the scope of the appended claims.

What is claimed is:

1. A drug delivery device comprising: an intraluminal stent; a biocompatible, nonerodible polymeric coating affixed to the intraluminal stent; and from about 64  $\mu$ g to about 197  $\mu$ g of rapamycin or a macrocyclic triene analog thereof that binds FKBP12 incorporated into the polymeric coating, wherein said device provides an in-stent late loss in diameter at 12 months following implantation in a human of less than about 0.5 mm, as measured by quantitative coronary angiography.

2. A drug delivery device according to claim 1 that provides an in-stent late loss in diameter at 12 months following implantation in a human of less than about 0.3 mm, as measured by quantitative coronary angiography.

3. A drug delivery device according to claim 1 or 2 that provides an in-stent diameter stenosis at 12 months following implantation in a human of less than about 22%, as measured by quantitative coronary angiography.

4. A drug delivery device according to claim 3 that provides an in-stent diameter stenosis at 12 months following implantation in a human of less than about 15%, as measured by quantitative coronary angiography.

5. A drug delivery device comprising: an intraluminal stent; a biocompatible, nonerodible polymeric coating affixed to the intraluminal stent; and from about 64  $\mu$ g to about 197  $\mu$ g of rapamycin or a macrocyclic triene analog thereof that binds FKBP12 incorporated into the polymeric coating, wherein said device provides a mean in-stent late loss in diameter in a human population at 12 months following implantation of less than about 0.5 mm, as measured by quantitative coronary angiography.

6. A drug delivery device according to claim 5 that provides a mean in-stent late loss in diameter in a human population at 12 months following implantation of less than about 0.3 mm, as measured by quantitative coronary angiography.

7. A drug delivery device according to claim 5 or 6 that provides a mean in-stent diameter stenosis in a human population at 12 months following implantation of less than about 22%, as measured by quantitative coronary angiography.

8. A drug delivery device according to claim 7 that provides a mean in-stent diameter stenosis in a human population at 12 months following implantation of less than about 15%, as measured by quantitative coronary angiography.

18

9. A method of inhibiting neointimal proliferation in a human coronary artery resulting from percutaneous transluminal coronary angioplasty comprising implanting in the lumen of said coronary artery a drug delivery device comprising: an intraluminal stent; a biocompatible, nonerodible polymeric coating affixed to the intraluminal stent; and from about 64  $\mu$ g to about 197  $\mu$ g of rapamycin or a macrocyclic triene analog thereof that binds FKBP12 incorporated into the polymeric coating, wherein said method provides an in-stent late loss in diameter at 12 months following implantation of less than about 0.5 mm, as measured by quantitative coronary angiography.

10. A method according to claim 9 that provides an in-stent late loss in diameter at 12 months following implantation of less than about 0.3 mm, as measured by quantitative coronary angiography.

11. A method according to claim 9 or 10 that provides an in-stent diameter stenosis at 12 months following implantation of less than about 22%, as measured by quantitative coronary angiography.

12. A method according to claim 11 that provides an in-stent diameter stenosis at 12 months following implantation of less than about 15%, as measured by quantitative coronary angiography.

13. A method of inhibiting neointimal proliferation in a coronary artery resulting from percutaneous transluminal coronary angioplasty comprising implanting in the lumen of said coronary artery a drug delivery device comprising: an intraluminal stent; a biocompatible, nonerodible polymeric coating affixed to the intraluminal stent; and from about 64  $\mu$ g to about 197  $\mu$ g of rapamycin or a macrocyclic triene analog thereof that binds FKBP12 incorporated into the polymeric coating, wherein said method provides a mean in-stent late loss in diameter in a human population at 12 months following implantation of less than about 0.5 mm, as measured by quantitative coronary angiography.

14. A method according to claim 13 that provides a mean in-stent late loss in diameter in a human population at 12 months following implantation of less than about 0.3 mm, as measured by quantitative coronary angiography.

15. A method according to claim 13 or 14 that provides a mean in-stent diameter stenosis in a human population at 12 months following implantation of less than about 22%, as measured by quantitative coronary angiography.

16. A method according to claim 15 that provides a mean in-stent diameter stenosis in a human population at 12 months following implantation of less than about 15%, as measured by quantitative coronary angiography.

17. The drug delivery device according to any one of claims 1, 2, 4 or 5 wherein said rapamycin or macrocyclic triene analog thereof is incorporated into the polymeric coating at a dose of from about 64  $\mu$ g to about 125  $\mu$ g.

18. The drug delivery device according to any one of claims 1, 2, 4 or 5 that releases a portion of said dose of rapamycin or a macrocyclic triene analog thereof at about six weeks following intraluminal implantation.

19. The drug delivery device according to any one of claims 1, 2, 4 or 5 wherein said rapamycin or macrocyclic triene analog thereof is incorporated into the polymeric coating at a dose of from about 2  $\mu$ g to about 30  $\mu$ g per millimeter of stent length.

20. The drug delivery device according to claim 19 wherein said rapamycin or macrocyclic triene analog thereof is incorporated into the polymeric coating at a dose of from about 3  $\mu$ g to about 13  $\mu$ g per millimeter of stent length.

A1256

US 7,300,662 B2

19

21. The drug delivery device according to claim 19 that releases a portion of said dose of rapamycin or a macrocyclic triene analog thereof at about six weeks following intraluminal implantation.

22. The method according to any one of claims 9, 10, 13 or 14, wherein said rapamycin or macrocyclic triene analog thereof is incorporated into the polymeric coating at a dose of from about 64  $\mu$ g to about 125  $\mu$ g.

23. The method according to any one of claims 9, 10, 13 or 14, wherein said rapamycin or macrocyclic triene analog thereof is incorporated into the polymeric coating at a dose of from about 2  $\mu$ g to about 30  $\mu$ g per millimeter of stent length.

20

24. The method according to any one of claims 9, 10, 13 or 14, wherein said rapamycin or macrocyclic triene analog thereof is incorporated into the polymeric coating at a dose of from about 3  $\mu$ g to about 13  $\mu$ g per millimeter of stent length.

25. The method according to any one of claims 9, 10, 13 or 14, wherein said drug delivery device releases a portion of said dose of rapamycin or a macrocyclic triene analog thereof at about six weeks following intraluminal implantation.

\* \* \* \* \*

JACC



## Journal of the American College of Cardiology/Contents in brief

OCTOBER 1995 VOLUME 26 NUMBER 4

## CLINICAL STUDIES

## ELECTROPHYSIOLOGY

- 843 Radiofrequency Catheter Ablation for Management of Symptomatic Ventricular Ectopic Activity Dennis Wei-Xi Zhu, James D. Maloney, Tony W. Simmons, Junichi Nitta, David M. Fitzgerald, Richard G. Trohman, Dirar S. Khoury, Walid Saliba, Karen M. Belco, Carlos Rizo-Patron, Sergio L. Pinski
- 850 Editorial Comment Radiofrequency Catheter Ablation of Benign Ventricular Ectopic Beats: A Therapy in Search of a Disease? Hein J. J. Wellens
- 852 Efficacy and Proarrhythmic Hazards of Pharmacologic Cardioversion of Atrial Fibrillation: Prospective Comparison of Sotalol Versus Quinidine Stefan H. Hohnloser, Andreas van de Loo, Frank Baedeker
- 859 Association Between QT Dispersion and Autonomic Dysfunction in Patients With Diabetes Mellitus Kevin Wei, Paul Dorian, David Newman, Anatoly Langer
- 864 High Prevalence of Antibodies Against Beta<sub>1</sub>- and Beta<sub>2</sub>-Adrenoceptors in Patients With Primary Electrical Cardiac Abnormalities Pablo A. Chiale, Mauricio B. Rosenbaum, Marcelo V. Elizari, Ake Hjalmarson, Yvonne Magnusson, Gerd Wallukat, Johan Hoebeke

## MYOCARDIAL ISCHEMIA

- 870 Evaluation of Myocardial Ischemia Using a Rest Metabolism/Stress Perfusion Protocol With Fluorine-18 Deoxyglucose/Technetium-99m MIBI and Dual-Isotope Simultaneous-Acquisition Single-Photon Emission Computed Tomography Martin P. Sandler, Searle Videlefsky, Dominique Delbecq, James A. Patton, Colin Meyerowitz, William H. Martin, Israel Ohana
- 879 Long-Term Prediction of Major Ischemic Events by Exercise Thallium-201 Single-Photon Emission Computed Tomography: Incremental Prognostic Value Compared With Clinical, Exercise Testing, Catheterization and Radiolabeled Angiographic Data Pierre-Yves Marie, Nicolas Danchin, Jean F. Durand, Luc Feldmann, Alain Grentzinger, Pierre Olivier, Gilles Karcher, Yves Juillière, Jean M. Virion, Daniel Beurrier, François Cherrier, Alain Bertrand
- 887 Prognostic Value of High Dose Dipyridamole Echocardiography in Patients With Chronic Coronary Artery Disease and Preserved Left Ventricular Function Claudio Coletta, Alfonso Galati, Gabriella Greco, Maurizio Burattini, Roberto Ricci, Alessandro Carunchio, Maria Stella Fera, Loretta Bordini, Vincenzo Ceci
- 895 Recurrence of Angina After Coronary Artery Bypass Surgery: Predictors and Prognosis (CASS Registry) Airlie A. C. Cameron, Kathryn B. Davis, William J. Rogers

## MYOCARDIAL INFARCTION

- 900 Prognostic Significance of Nonfatal Reinfarction During 3-Year Follow-Up: Results of the Thrombolysis in Myocardial Infarction (TIMI) Phase II Clinical Trial Håvard S. Mueller, Sandra A. Forman, Mark A. Menegus, Lawrence S. Cohen, Genell L. Knatterud, Eugene Braunwald, for the TIMI Investigators
- 908 Stress Echocardiographic Results Predict Risk of Reinfarction Early After Uncomplicated Acute Myocardial Infarction: Large-Scale Multicenter Study Eugenio Picano, Alessandro Pingitore, Rosa Sicari, Giovanni Minardi, Nicola Gandolfo, Giovanni Seveso, Francesco Chiarella, Leonardo Bolognese, Giacomo Chiarandà, Maria Grazia Schiavo, Mario Previtali, Franca Margaria, Ornella Magnaia, Federico Bianchi, Salvatore Pirelli, Silva Severi, Mauro Raciti, Patricia Landi, Cristina Vassalle, Maria José Bento de Sousa, Luis Felipe de Moura Duarte, on Behalf of the Echo Persantine International Cooperative (EPIC) Study Group
- 914 Cost-Effectiveness of Captopril Therapy After Myocardial Infarction Joel Tsevat, Daniela Duke, Les Goldman, Marc A. Pfeffer, Gervasio A. Lamas, Jane R. Soukup, Karen M. Kuntz, Thomas H. Lee
- 920 Editorial Comment Cost-Effectiveness Analysis and Clinical Practice Thomas S. Rector, Gary S. Francis
- 922 INTERVENTIONAL CARDIOLOGY  
Peripheral Vascular Complications in the Coronary Angioplasty Versus Excisional Atherectomy Trial (CAVEAT-I) Nowamugbe A. Omoigui, Robert M. Califf, Karen Pieper, Gordon Keeler, Mary Ann O'Hanorian, Lisa G. Berdan, Daniel B. Mark, J. David Talley, Eric J. Topol, for the CAVEAT-I Investigators
- 931 Development and Validation of a Simplified Predictive Index for Major Complications in Contemporary Percutaneous Transluminal Coronary Angioplasty Practice Stephen E. Kimmel, Jesse A. Berlin, Brian L. Strom, Warren K. Laskey, for the Registry Committee of the Society for Cardiac Angiography and Interventions
- 939 Immediate and Late Outcome of Excimer Laser and Balloon Coronary Angioplasty: A Quantitative Angiographic Comparison Based on Matched Lesions Sipke Strikwerda, Eline Montauban van Swijndregt, David P. Foley, Eric Boersma, Victor A. Umans, Rein Melker, Patrick W. Serruys
- 947 The Subcutaneous Heparin and Angioplasty Restenosis Prevention (SHARP) Trial: Results of a Multicenter Randomized Trial Investigating the Effects of High Dose Unfractionated Heparin on Angiographic Restenosis and Clinical Outcome Michael J. Brack, Simon Ray, Anoop Chauhan, Johnathan Fox, Peter J. B. Hubner, Peter Schofield, Alex Harley, Anthony H. Gershlick, on Behalf of the SHARP Trial Investigators
- 955 Percutaneous Revascularization of Ostial Saphenous Vein Graft Stenoses Alaa E. Abdel-Meguid, Patrick L. Whitlow, Conrad

Journal of the American College of Cardiology (ISSN 0735-1097) is issued monthly, except semimonthly in March and November, in two indexed volumes per year by Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010. Printed in USA at 300 Cadmus Lane, Eason, MD 21601-0969. Subscription prices per year: Institution, \$180.00; individual, \$117.00; interns, residents, and allied health professionals, \$72.00. Outside USA, add \$25.00 for surface postage and handling. For air delivery to USA, Canada and Mexico, add \$143.00; to Europe, \$198.00 (via surface air-lift); to Japan, \$265.00; and to rest of world \$355.00. Second-class postage paid at New York, NY and at additional mailing offices. Postmaster: Send address changes to Journal of the American College of Cardiology, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010.



0735-1097(199510)26:4;1-V

CORD086409

A1258

## The Subcutaneous Heparin and Angioplasty Restenosis Prevention (SHARP) Trial

### Results of a Multicenter Randomized Trial Investigating the Effects of High Dose Unfractionated Heparin on Angiographic Restenosis and Clinical Outcome

MICHAEL J. BRACK, MD, SIMON RAY, MD,\* ANOOP CHAUHAN, MD,† JOHNATHAN FOX, MRCP,\* PETER J. B. HUBNER, FRCP,‡ PETER SCHOFIELD, FRCP,† ALEX HARLEY, FRCP,\* ANTHONY H. GERSHLICK, FRCP,‡ ON BEHALF OF THE SHARP TRIAL INVESTIGATORS

Blackpool, Leicester, Liverpool and Cambridge, England, United Kingdom

**Objectives.** We sought to determine whether 12,500 IU of unfractionated heparin given subcutaneously twice daily for 4 months after percutaneous transluminal coronary angioplasty beneficially influences the subsequent rate of angiographic restenosis and the incidence of clinical events.

**Background.** Heparin has been shown to exhibit powerful antiproliferative effects against smooth muscle cells in several animal models.

**Methods.** A randomized trial with blinded data analysis was undertaken to assess the effect of unfractionated subcutaneous heparin on angiographic restenosis after coronary angioplasty. After successful angioplasty, patients were randomized to receive no heparin or 12,500 IU of heparin given subcutaneously twice daily for 4 months. Quantitative coronary angiography was performed before angioplasty, immediately after angioplasty and at follow-up ("early" [before 4 months] or electively [at 4 months]).

**Results.** The study group comprised 339 patients, 167 randomly

assigned to receive heparin, 172 to receive no heparin. Repeat cardiac catheterization was performed in 90% of randomized patients. At early and elective restudy (mean 4.2 months), the mean  $\pm$  SD difference in minimal lumen diameter between the postangioplasty and follow-up measurement was  $-0.55 \pm 0.58$  mm for the no heparin group and  $-0.43 \pm 0.59$  mm for the heparin group ( $p = \text{NS}$ ). Clinical events during the follow-up period did not differ significantly between groups: fatal myocardial infarction (1 patient in each group), coronary bypass grafting (5 patients in each group), repeat angioplasty (12 in the no heparin, 6 in the heparin group), angina at 4-month assessment (33% in the no heparin, 32% in the heparin group).

**Conclusions.** Long-term treatment with high dose subcutaneous heparin (12,500 IU twice daily) for 4 months did not favorably influence angiographic or clinical outcome after coronary angioplasty.

(*J Am Coll Cardiol* 1995;26:947-54)

Percutaneous transluminal coronary angioplasty is a widely accepted form of treatment for symptomatic coronary artery disease. The late restenosis rate of 30% to 60% (1-5) continues to limit the longer term benefit of the procedure.

The restenotic process is characterized by elastic recoil (up to 40%) and myointimal hyperplasia (6,7), which is initiated by migration of medial smooth muscle cells from the media to the intima (8), with proliferation there after phenotypic modulation of the migrating cells. Subsequent extracellular matrix makes up the bulk of the later (>1 month) restenotic lesion.

From the Blackpool Victoria Hospital, Blackpool; \*Cardiothoracic Centre, Liverpool; †Papworth Hospital, Cambridge; and ‡Department of Cardiology, Glenfield General Hospital, Leicester, England, United Kingdom. This project was supported by Project Grant 91/18 from The British Heart Foundation, London, England, United Kingdom. Heparin was supplied by CP Pharmaceuticals Ltd., Croyd, Wales, United Kingdom.

Manuscript received October 21, 1994; revised manuscript received April 7, 1995; accepted May 22, 1995.

Address for correspondence: Dr. Anthony H. Gershlick, Department of Cardiology, Glenfield General Hospital, Groby Road, Leicester LE3 9QP, England, United Kingdom.

Experimental work in animal models (9-15) and from in vitro studies (16) suggests that heparin may inhibit changes in intimal smooth muscle cells and therefore influence the restenotic process.

In a large North American (enoxaparin) trial (17), low molecular weight heparin was shown to have no effect on angiographic restenosis or clinical events. The present study, referred to as the Subcutaneous Heparin and Angioplasty Restenosis Prevention (SHARP) trial, was performed to assess the value of unfractionated calcium heparin in preventing restenosis after coronary angioplasty.

## Methods

**Study patients.** All symptomatic patients who had undergone successful coronary angioplasty (defined as visually assessed diameter stenosis <50% after angioplasty) for an angiographically proved narrowing in one or more coronary



arteries were considered for entry into the trial in three participating regional centers (Leicester, Liverpool and Papworth, England, United Kingdom). Patients with restenotic lesions, chronic occlusions or conduit lesions were not included. Ethical approval for the trial was granted by the appropriate committees at each participating center.

**Angioplasty procedure and follow-up angiography.** All angioplasty procedures were performed by the femoral route with use of steerable, movable guide wire systems. Aspirin (300 mg) was given to all patients before the procedure; the use of calcium channel antagonists or beta-adrenergic blocking agents, or both, was determined by the physician in charge of the patient. After placement of the femoral artery sheath all patients received a bolus of 10,000 IU of intravenous heparin. During prolonged procedures additional heparin (5,000 IU) was given after 1 h. After the procedure and immediately on the patient's return to the ward, intravenous administration of heparin was begun at an initial dose of 1,000 IU/h. The dose of heparin was adjusted to maintain the activated partial thromboplastin time ratio to 2.0 to 3.0:1 for up to 24 h. Balloon type and size, inflation pressure and duration of inflation were selected by each operator.

Three coronary angiograms were obtained from each patient: one immediately before and one immediately after coronary angioplasty and one at follow-up (either elective [at 4 months] or "early," if clinically indicated).

**Data acquisition.** Quantitative analysis was performed with use of the CardioTrace system, whose results have been shown (18) to correlate well with those of the Coronary Angiography Analysis System (CAAS) using phantoms. Essentially, orthogonal views of each lesion were taken and the exact positioning of the X-ray tube and table height noted. The angiograms obtained before and after angioplasty and at follow-up were taken in the same views. Before each angiographic sequence the contrast-free catheter was filmed and at the end of the procedure the distal 20 cm was cut and saved for subsequent micrometer measurement and calibration of the analysis system. Before the postangioplasty angiogram the guide wire was removed to avoid interference with automated edge detection. Lesion characteristics were also assessed qualitatively.

All angiographic analysis was performed at the core laboratory by a single operator who had no knowledge of the patient's group. Postangioplasty angiograms were examined for dissection defined according to the modified National Heart, Lung, and Blood Institute criteria. Intracoronary thrombus was defined as the presence of a filling defect within the lumen, surrounded by contrast material seen in multiple projections in the absence of calcium within the filling defect, the persistence of contrast material within the lumen or a visible embolization of intraluminal material downstream.

The absolute values of stenosis minimal lumen diameter and reference diameter were measured by using the catheter tip as a scaling device. The reference measurement was taken from the "normal" coronary segment at the time of stenosis measurement to allow for change in the reference artery with

time (19). The CardioTrace automatically adjusts for the catheter size as measured by the micrometer gauge.

**Treatment allocation.** After successful coronary angioplasty and on the patient's return to the ward, we sought the patient's consent to be included in this trial. Consent was sought at this time because this was a trial of successful angioplasty and subsequent restenosis. A total of 339 patients met all inclusion and exclusion criteria and gave informed written consent. A computer-generated program randomly assigned patients to receive either no heparin (no injection) or unfractionated subcutaneous heparin (12,500 IU twice daily for 4 months).

Patients randomized to the heparin group were given the injections 2 h after femoral sheath withdrawal. All patients were discharged receiving aspirin (75 to 300 mg daily) and calcium channel antagonists, nitrates and beta-adrenergic blocking agents according to operator choice.

**Follow-up evaluation.** Patients receiving heparin treatment had a peak plasma heparin assay performed 1 week after discharge and a weekly platelet count for the 1st 4 weeks of therapy. Anticoagulation levels were not measured, because this is the level of heparin given long-term and safely in other groups of patients.

A syringe count was performed at 2 and 4 months. If patients reported return of their angina within the 4-month follow-up period, coronary angiography was performed if the physician deemed it clinically indicated. Even if there was no evidence of restenosis patients were not required to undergo further catheterization at 4 months.


On admission for the 4-month elective catheterization, patients underwent a standard treadmill exercise test (Bruce protocol). End points for the test were the occurrence of anginal chest pain, decrease in systolic blood pressure, significant arrhythmia, or ST segment depression >1 mm 80 ms after the J point. Clinical status was assessed during the in-hospital admission for repeat catheterization.

**End points.** The primary end point of the SHARP trial was the within-patient change in minimal lumen diameter as determined by quantitative angiography. Postangioplasty values were obtained from the last postangioplasty angiogram after withdrawal of the guide wire. For patients admitted early who underwent repeat angioplasty, the angiogram preceding repeat balloon dilation was considered the follow-up angiogram and was recorded in exactly the same views used in the angiograms previously obtained immediately before and after angioplasty. For all other patients the planned follow-up angiogram was used.

Secondary end points were death, myocardial infarction, need for coronary artery bypass surgery and repeat coronary angioplasty. The presence and severity of angina at follow-up (early or elective) were assessed according to the Canadian Cardiovascular Society classification.

**Statistical analysis.** Calculations for the power of any restenosis trial are based on an arbitrary assessment of the biologic goal balanced by the side effects of the medication being tested. In an attempt to objectively determine the sort of

Figure 1. Patient flow from time of randomization to follow-up for the no heparin and heparin groups. "Early" and elective = cardiac catheterization performed, respectively, before 4 months and at the scheduled 4-month follow-up time.



effect we might achieve with heparin, we tested, in a pretrial study, the plasma of patients given the study dose of heparin (12,500 IU twice daily) against human smooth muscle cell growth in vitro. We found a 35% reduction in smooth muscle cell growth over 10 days (from  $6.9 \times 10^4$  to  $4.5 \times 10^4$ ) when these cells were exposed to blood from patients treated with heparin (mean level 0.3 IU/ml). Because this level of reduction is approximately equal to the arbitrarily used 33% reduction deemed important by other trialists, we used it as the aim of the treatment arm. On the basis of this 35% reduction and the results of Nobuyoshi et al. (20) and Serruys et al. (3), we estimated that the mean  $\pm$  SD loss in minimal lumen diameter over a 4- to 6-month period in control patients would be  $-0.50 \pm 0.5$  mm. Assuming that diameters are normally distributed (21), we calculated that to show a reduction (the anticipated loss of lumen diameter by 35% in the heparin-treated group), 129 patients with complete angiographic follow-up were required in each group to give a power of 80% at an alpha level of 0.05. To allow for medical withdrawal of patients and patients defaulting from the trial, we aimed to recruit 340 patients. A total of 339 patients who had had successful angioplasty were recruited.

## Results

**Patient numbers.** From February 1990 to the end of June 1993, 370 patients who had undergone successful coronary angioplasty were approached for participation in the SHARP trial. A total of 339 patients (92%) consented to participate; of these, 167 patients (189 lesions) were randomly assigned to receive subcutaneous heparin (12,500 U twice daily) and 172 patients (188 lesions) to receive no heparin. During the trial period 40 patients defaulted from the trial (27 in the heparin and 13 in the no heparin group); 39 patients required "early" cardiac catheterization (23 in the no heparin and 16 in the heparin group). Figure 1 shows the patient flow from the time of randomization to follow-up.

**Baseline characteristics and clinical follow-up.** Tables 1 and 2, respectively, show the demographic data for the two groups after default and lesion characteristics before and after default. The baseline characteristics of both groups were well matched. Clinical follow-up data were available from all patients not defaulting from the trial. During the course of the

Table 1. Patient Characteristics of the No Heparin and Heparin Groups After Default of 40 Patients

	No Heparin Group (129 pts, 174 lesions)	Heparin Group (140 pts, 161 lesions)
Men (%)	82	80
Age (yr) (mean $\pm$ SD)	56.2 $\pm$ 7	56.7 $\pm$ 9
Smoker	48	56
Diabetes	11	7
Hypertension	31	26
Hyperlipidemia	29	26
Previous MI	71	64
Previous CABG	6	3
Previous PTCA	3	4
Drug therapy		
Aspirin	122	109
Calcium antagonists	106	101
Nitrates	104	91
Beta-blockers	102	96
Angina grade		
0	0	0
1	4	3
2	81	73
3	55	49
4	19	15
Duration of angina (mo)		
<6	40	41
6-12	53	42
12-24	21	18
>24	45	59

Unless otherwise indicated, data are expressed as number of patients. CABG = coronary artery bypass grafting; MI = myocardial infarction; PTCA = percutaneous transluminal coronary angioplasty; pts = patients.

Table 2. Vessel and Lesion Characteristics of the Heparin and No Heparin Groups Before and After Definitive

	No Heparin Group		Heparin Group	
	Pre	Post	Pre	Post
Vessels	188	174	189	161
Vessel dilated				
LAD	100	95	100	87
RCA	46	42	49	40
LCx	42	37	40	34
Stents dilated (no.)				
1	160	150	150	124
2	12	10	15	14
3	0	0	3	3
4	1	1	0	0
Lesion characteristic				
A	64	59	67	57
B	97	91	104	92
C	27	24	18	12
Dissections (no.)	38	33	34	27
Dissection grade				
A	14	12	10	9
B	13	13	12	9
C	9	7	10	8
D	2	1	2	1
E	0	0	0	0
Reference artery/ balloon ratio	0.96/0.17	0.17/0.2	0.90/0.14	1.0/0.2

Unless otherwise indicated, data are expressed as number of vessels. LAD = left anterior descending coronary artery; LCx = left circumflex coronary artery; Pre (Post) = before (after) definitive; RCA = right coronary artery.

study two patients (one patient in each group) died, both of myocardial infarction. Ten patients (5 in each group) underwent coronary bypass surgery and 18 (12 in the no heparin and 6 in the heparin group) had repeat coronary angioplasty. At the 4-month assessment, 33% of patients in the no heparin and 32% of those in the heparin group reported recurrent angina.

**Angiographic analysis.** During the study period, 39 patients underwent "early" cardiac catheterization because anginal symptoms had recurred (23 patients in the no heparin group and 16 in the heparin group). Table 3 shows the reference diameters, minimal lumen diameters, restenosis rates and clinical outcome of the patients undergoing early catheterization. The minimal lumen diameters before and after coronary angioplasty and at follow-up were not significantly different at any time between the two patient groups. As assessed by a definition of "loss of 50% of the absolute gain," 19 patients in the no heparin and 9 in the heparin group had restenosis, (chi-square test,  $p = 0.15$ , odds ratio 0.27, 95% confidence interval [CI] -0.06 to 1.2). There were no significant differences in restenosis rates or changes in minimal lumen diameter between patients with early or elective follow-up angiography.

Figure 2 shows the quantitative angiographic details of the two groups at elective follow-up. For those patients undergoing elective repeat coronary angiography, the difference between the minimal lumen diameter at follow-up and immediately

Table 3. Minimal Lumen Diameters and Clinical Outcome of Patients Undergoing "Early" (&lt;4 mo) Repeat Cardiac Catheterization

	No Heparin Group (n = 23)	Heparin Group (n = 16)
Reference diameter (mm)		
Pre PTCA	2.81 ± 0.56	2.71 ± 0.50
Post PTCA	2.86 ± 0.49	2.69 ± 0.55
Follow-up	2.76 ± 0.46	2.76 ± 0.55
Minimal lumen diameter (mm)		
Pre PTCA	0.68 ± 0.42	0.78 ± 0.31
Post PTCA	1.78 ± 0.42	1.76 ± 0.48
Follow-up	0.96 ± 0.71	1.18 ± 0.59
% diameter reduction		
Pre PTCA	75.3 ± 15	72.3 ± 10
Post PTCA	37.1 ± 14	34.5 ± 11
Follow-up	64.6 ± 25	57.3 ± 20
Angina grade (CCS class)		
Mean	2.6	2.6
Range	2-4	1-4
Repeat catheterization at (wk)		
Mean	8.5	7.5
Range	1-14	0.5-12
Repeat PTCA	12	6
CABG	5	5
Medical therapy	6	5

Unless otherwise indicated, data are presented as number of patients or mean value ± SD. CCS class = Canadian Cardiovascular Society classification; other abbreviations as in Tables 1 and 2.


after angioplasty was  $-0.49 \pm 0.57$  mm in the no heparin group and  $-0.41 \pm 0.57$  mm in the heparin group. The 0.88-mm difference—that is, the treatment effect—was not significant ( $p = 0.22$ , 95% CI -0.05 to 0.22). As assessed by a categorical definition of loss of 50% of the absolute gain, the restenosis rate was 46% in the no heparin and 39% in the heparin group (chi-square test,  $p = 0.32$ , odds ratio 0.75, 95% CI 0.46 to 1.2). For all patients (those with early and elective restudy), the loss at follow-up in minimal lumen diameter was  $-0.55 \pm 0.58$  mm for the no heparin and  $-0.43 \pm 0.59$  mm in the heparin group (Table 4). The overall treatment effect was 0.12 mm ( $p = 0.07$ , 95% CI -0.004 to -0.25 mm), which corresponded to a restenosis rate of 51% in the no heparin and 41% in the heparin group ( $p = 0.09$ ).

Figures 3 and 4 represent a cumulative frequency curve of the minimal lumen diameter and a plot of relative loss versus relative gain for the two groups.

**Treadmill exercise tests.** At follow-up 85% of patients in the no heparin and 86% of those in the heparin group underwent a standard treadmill exercise test (Bruce protocol). There were no differences between groups in the number of patients reporting pain or having significant ST changes during the test.

**Acceptability of heparin treatment.** The number of patients randomized into the study represents 92% of the patients approached. The main reason for nonenrollment was unwillingness to self administer heparin. Indeed, the major

Figure 2. Reference diameters, minimal lumen diameters before (Pre) and immediately after (Post) percutaneous transluminal coronary angioplasty (PTCA) and at follow-up for the two groups. Acute gain and late loss are also shown.



reason for default was the inconvenient nature of the treatment. Three patients were withdrawn from the study as a result of pruritic swellings at the site of the injections that resolved on discontinuation of treatment. No cases of thrombocytopenia were reported and the occurrence of osteoporosis in one patient in each group is of doubtful significance. Apart from two instances of bleeding at the femoral puncture site, no patients reported spontaneous bruising or troublesome hematoma formation after minor trauma. The treatment, although onerous, would appear to be safe and generally well tolerated.

Table 4. Absolute Measurements of Reference Diameters and Obstruction Diameters for All Patients Undergoing Repeat Catheterization

	No Heparin Group (159 pts, 174 lesions)	Heparin Group (140 pts, 161 lesions)
Reference diameter (mm)		
Pre PTCA	2.82 ± 0.54	2.87 ± 0.54
Post PTCA	2.81 ± 0.50	2.87 ± 0.51
Follow-up	2.81 ± 0.50	2.87 ± 0.52
Minimal lumen diameter (mm)		
Pre PTCA	0.83 ± 0.37	0.81 ± 0.34
Post PTCA	1.91 ± 0.46	1.88 ± 0.48
Follow-up	1.36 ± 0.63	1.45 ± 0.64
Acute gain (mm)	1.08 ± 0.48	1.07 ± 0.49
Relative gain	0.39 ± 0.18	0.38 ± 0.15
Late loss (mm)	0.55 ± 0.58	0.43 ± 0.59
Relative loss	0.20 ± 0.22	0.15 ± 0.21
Net gain index	0.19 ± 0.22	0.22 ± 0.20
Percent stenosis		
Pre PTCA	71.5 ± 12%	71.9 ± 10%
Post PTCA	31.6 ± 13%	35.4 ± 12%
Follow-up	52.0 ± 21%	50.0 ± 19%
Restenosis rate (loss >50% absolute gain)		
Patients	51%	41%
Vessels	48%	39%

Unless otherwise indicated, data are expressed as mean value ± SD. Abbreviations as in Tables 1 and 2.

## Discussion

Although the effects of heparin on smooth muscle cell proliferation have been known for some time, the precise mode of action is unclear. Heparin binds to smooth muscle cells and is taken up by both receptor-mediated and slower endocytotic pathways (22). Studies utilizing ornithine decarboxylase activity and tritiated thymidine uptake suggest that heparin acts either at the late G1 or early S phase of the cell cycle, with consequent reduction in synthesis of deoxyribonucleic acid and ribonucleic acid (23). Heparin may also have secondary effects due to a reduction in expression of growth factor receptors (24), or enhanced action of growth inhibitory factors (25) such as transforming growth factor B.

There are few studies in humans addressing the effect of heparin on restenosis after coronary angioplasty. The trial by Ellis et al. (26), randomly assigned 416 patients to receive either intravenous heparin or dextrose for 18 to 24 h after coronary angioplasty. The restenosis rates were 41.2% in the heparin group and 36.7% in the control group when assessed at 6 months (restenosis defined as >50% diameter stenosis at follow-up). Similar results were obtained by Saenz et al. (27), although their study had considerably fewer patients. For similar reasons little meaningful data could be derived from the study by Lehmann et al. (28), which was terminated early on ethical grounds because 82% of the 23 patients at follow-up had restenosis. More data are available from the Enoxaparin study (17) involving 465 patients who were randomized to low molecular weight heparin (30 mg subcutaneously daily) or placebo for 1 month. As assessed from clinical end points, the restenosis rate was 51% in the treatment group and 49% in the control group. There was no significant difference in the change in minimal lumen diameter at follow-up (0.49 and 0.5 mm, respectively, for the control and the treatment groups).

There has been some debate as to whether the commercially available low molecular weight heparin contains all of the antiproliferative effects of the subfractions outlined in the publications by Guyton et al. (10) and Hoover et al. (29).





Figure 3. Cumulative frequency curves of minimal lumen diameters for the heparin (triangles) and no heparin (circles) groups before and immediately after coronary angioplasty and at follow-up.

Commercially available low molecular weight heparin may not be the same subtraction as that used in experimental models. For this reason we chose to examine the effects of unfractionated heparin, knowing also that any retained antithrombotic effects of the unfractionated heparin may also be beneficial if thrombus formation is another smooth muscle cell stimulus (30). We chose to give the heparin for a total of 4 months because data suggest that 90% of restenosis if it is to occur will occur within this time. Although the process of smooth muscle cell proliferation takes place in the first few days, we chose not to risk merely delaying the process by limiting the time of heparin administration to this period.

Despite the advantages of using this relatively high dose of unfractionated heparin given over the known course of the restenotic process, we have shown that this treatment failed to beneficially influence either the angiographic outcome (amount of intimal hyperplasia) or the clinical outcome. There may have been several reasons for these findings.

Figure 4. Plot of relative loss versus relative gain for the heparin (plus signs) and no heparin (circles) groups.



**Study power.** Currently the power of any trial is determined by the effect one wants to achieve (31). Thus a 10% reduction requires larger numbers than if a 50% reduction is the aim. The use by other investigators of a 33% reduction (resulting in trial numbers of 230 patients/group) is as arbitrary as any other present reduction. Our choice was based on *in vitro* studies. However, for the secondary end points (clinical outcome) the study power was too small.

**Heparin dose.** The dose of heparin selected for this trial was based on both ethical and pharmacologic considerations. At the time of the study's inception, the largest available dose of "ready to use" heparin was 12,500 IU. Our dose of 25,000 IU of heparin daily approaches doses that could result in a significant risk that patients would develop osteoporosis (32). Additionally, the statements of patients interviewed before the trial, but not part of it, indicated that asking patients to self-inject this drug more than twice daily would be unacceptable.

From our own laboratory work using cultured human smooth muscle cells we found that concentrations of heparin that could be achieved by subcutaneous injection of 12,500 IU (i.e., 0.3 IU/ml), significantly inhibited smooth muscle cell proliferation. Castellotti et al. (33) have shown that 1 to 5  $\mu$ g of heparin added to growth media could produce inhibition of smooth muscle cells in culture. However, *in vivo* work suggests that higher doses are required to produce a significant inhibition of smooth muscle cell proliferation. Clowes and Karnovsky (9), using the rat carotid model, reported that 50 to 100 IU/kg body weight per h was needed to produce significant inhibition of cell proliferation. Likewise, using hypercholesterolemic rabbits, Currier et al. (13) showed that 10 mg/kg per day of enoxaparin could reduce the restenosis rate in iliac arteries, but human studies (17) using 30 mg/day showed no benefit.

Thus, although we used the largest ethically acceptable dose, this dose may have been too small to locally inhibit smooth muscle cell proliferation. There has been a suggestion, perhaps a justification of negative results, that humans are

more sensitive than animals to many of the drugs tried systemically, and thus the dosage issue is not relevant. However, in the light of our results, it is reasonable to conclude that human vascular smooth muscle cells are no more sensitive than animal smooth muscle cells to the effects of heparin; otherwise, some beneficial effect ought to have been observed with this smaller dose.

**Conclusions.** The failure of unfractionated heparin to show any benefit in humans despite encouraging results in animal models is similar to the outcome of trials with cilazapril (34), enoxaparin (17) and ketanserlin (35). The ability of heparin to inhibit vascular smooth muscle cell growth *in vitro* and in animal models of vascular injury is well documented. It appears not to have the therapeutic efficacy to reduce restenosis after angioplasty in humans. The published results of intravenous short-term administration (24 to 48 h) of heparin and the results of medium-term (1 month) of low molecular weight heparin also show no benefit. The reasons for these findings have been addressed by other trialists studying other drugs, namely the discrepancy between systemic doses that work in animals and those that can be given to humans. The argument that smooth muscle cells or the mechanisms stimulating them are more sensitive in humans appears not to hold. Although we undertook a study with heparin given in as large a dose that was believed to be safe and ethical, it is clear that attention will need to be turned to methods of local delivery that will permit administration of doses that affect the process without adversely affecting the patient. Perhaps before heparin is finally rejected as a "magic bullet" for the prevention of restenosis, clinical trials should be conducted that match animal studies with continuous infusions or by local targeted delivery.

We are grateful for the assistance in the design of this study and the following persons for providing patients: Professor D. P. De Bono, Dr. J. D. Skehan, Dr. N. S. Samani, Mrs. L. Armstrong (M.S.O.), Glenfield General Hospital, Leicester; Dr. R. A. Perry, Dr. W. C. Morrison, Dr. D. R. Ramsdale, Dr. R. G. Charles, Dr. G. Epstein, Dr. S. Ray, Sister R. Carey, Cardiothoracic Centre, Liverpool; and Dr. M. C. Petch, Dr. D. L. Stone, Dr. L. M. Shapiro, Papworth Hospital, Cambridge, England, United Kingdom.

## Appendix

### Definitions of Lesion Characteristics and Dissection Grades (Table 2)

#### Lesion Characteristics

- Type A:** Small radiolucent area within the lumen of the vessel disappearing with the passage of the contrast material.  
**Type B:** Contrast extravasation parallel to the lumen of the vessel disappearing with the passage of the contrast material.  
**Type C:** Contrast extravasation persisting after passage of the contrast material.  
**Type D:** Spiral-shaped filling defect with normal (D1) or delayed (D2) antegrade flow.

**Type E:** Persistent lumen filling defect with delayed runoff of the contrast material in the distal vessel.

**Type F:** Filling defect accompanied by total occlusion.

#### Dissection Characteristics

- Type A:** Discrete, concentric, readily accessible, smooth contour, nonangulated segment, no calcification, not occlusive, not ostial, no major branch involvement, absence of thrombus.  
**Type B:** Tubular, eccentric, moderate tortuosity, moderately angulated segment, irregular contour, moderate calcification, total occlusion <3 months, ostial site, bifurcation lesion, thrombus present.  
**Type C:** Diffuse, very tortuous, markedly angulated, total occlusion >3 months, inability to protect side branch.

## References

- Holmes DR, Vlietstra RE, Smith HC, et al. Restenosis after percutaneous transluminal coronary angioplasty (PTCA): a report from the PTCA Registry of the National Heart, Lung, and Blood Institute. *Am J Cardiol* 1984;53:77C-81C.
- McBride W, Lange RA, Hillis LD. Restenosis after successful coronary angioplasty. *N Engl J Med* 1988;318:1334-7.
- Serruys FW, Luijckx HE, Beatt EJ, et al. Incidence of restenosis after successful coronary angioplasty: a time related phenomenon. A quantitative angiographic study in 342 consecutive patients at 1, 2, 3 and 4 months. *Circulation* 1988;77:361-71.
- Kuntz RE, Safian RD, Levine MJ, Reis GL, Dover DJ, Bain DS. Novel approach to the analysis of restenosis after the use of three new coronary devices. *J Am Coll Cardiol* 1992;19:1493-9.
- Hermans WR, Rensing BJ, Strauss BH, Serruys FW. Prevention of restenosis after percutaneous transluminal coronary angioplasty: the search for a "magic bullet." *Am Heart J* 1991;122:171-86.
- Austin GE, Ratcliffe NB, Hollman J, Tabei S, Phillips DF. Intimal proliferation of smooth muscle cells as an explanation for recurrent coronary artery stenosis after percutaneous transluminal coronary angioplasty. *J Am Coll Cardiol* 1985;6:369-75.
- Nobuuchi M, Kimura T, Ohishi H, et al. Restenosis after percutaneous transluminal coronary angioplasty: pathologic observations in 20 patients. *J Am Coll Cardiol* 1991;17:433-9.
- Ohara T, Aizawa S, Aizawa S. Ultrastructural study of proliferating smooth muscle cells: the site of PTCA as an explanation for restenosis [abstract]. *Circulation* 1988;78 Suppl II:II-290.
- Clowes AW, Karnovsky MJ. Suppression by heparin of smooth muscle cell proliferation in injured arteries. *Nature* 1977;266:525-6.
- Guyton JR, Rosenberg RD, Clowes AW, Karnovsky MJ. Inhibition of rat arterial smooth muscle cell proliferation by heparin. *In vivo* studies with anticoagulant and nonanticoagulant heparin. *Circ Res* 1980;46:623-34.
- Clowes AW, Clowes MM. Kinetics of cellular proliferation after arterial injury. IV. Heparin inhibits rat smooth muscle cell mitogenesis and migration. *Circ Res* 1986;58:839-45.
- Edelman ER, Adams DH, Karnovsky MJ. Effect of controlled adventitial heparin delivery on smooth muscle cell proliferation following endothelial injury. *Proc Natl Acad Sci U S A* 1990;87:3773-7.
- Currier JW, Pow TK, Hauschild CC, Minahan AC, Faxon DP. Low molecular weight heparin (enoxaparin) reduces restenosis after iliac angioplasty in the hypercholesterolemic rabbit. *J Am Coll Cardiol* 1991;17:1158-25B.
- Cimpe LW, Gertz SD, Haber HL, et al. Effect of chronic subcutaneous or intramuscular administration of heparin on femoral artery stenosis after balloon angioplasty in hypercholesterolemic rabbits. A quantitative angiographic and histopathological study. *Circulation* 1992;86:1536-46.
- Gordon JB, Berk BC, Bettmann MA, Selwyn AP, Renke H, Alexander RW. Vascular smooth muscle cell proliferation following balloon injury is synergistically inhibited by low molecular weight heparin and hydrocortisone [abstract]. *Circulation* 1987;76 Suppl IV:IV-213.
- Hoover RL, Rosenberg R, Haering W, Karnovsky MJ. Inhibition of rat

- arterial smooth muscle cell proliferation by heparin. II. in vitro studies. *Circ Res* 1980;47:578-83.
17. Faxon D, Spiro TE, Minor S, et al. Low molecular weight heparin in prevention of restenosis after angioplasty. Results of Enoxaparin Restenosis Cera Trial. *Circulation* 1994;90:932-14.
  18. Alvarez LG, Jackson SA, Berry A, Elmhorn EJ. Evaluation of a personal computer-based quantitative coronary analysis system for rapid assessment of coronary stenoses. *Am Heart J* 1992;123:1500-10.
  19. Beart KJ, Luijten HE, de Feyter PJ, van der Brand M, Reiber JHC, Serruys PW. Change in diameter of coronary artery segments adjacent to stenosis after percutaneous transluminal coronary angioplasty: failure of percent diameter stenosis measurements to reflect morphologic changes induced by balloon dilatation. *J Am Coll Cardiol* 1988;12:615-23.
  20. Nobuyoshi M, Kimura T, Nosaka H, et al. Restenosis after successful percutaneous transluminal coronary angioplasty: serial angiographic follow-up of 299 patients. *J Am Coll Cardiol* 1988;12:616-23.
  21. Reusing BJ, Hermans WR, Deckers JW, de Feyter PJ, Tijssen JGP, Serruys PW. Luminal narrowing after percutaneous transluminal coronary angioplasty follows a near Gaussian distribution: a quantitative angiographic study in 1445 successfully dilated lesions. *J Am Coll Cardiol* 1992;19:935-45.
  22. Castellor JJ, Wong K, Herman B, et al. Binding and internalization of heparin by vascular smooth muscle cells. *J Cell Physiol* 1985;124:13-20.
  23. Majesky MW, Schwartz SM, Clowes MM, Clowes AW. Heparin regulates smooth muscle S phase entry in injured rat carotid artery. *Circ Res* 1987;61:296-300.
  24. Reilly CF, Frize LMS, Rosenberg RD. Antiproliferative effects of heparin on vascular smooth muscle cells are reversed by epidermal growth factor. *J Cell Physiol* 1987;131:149-57.
  25. McCaffrey TA, Falcone DJ, Brayton CE, Agarwal LA, Welt FG, Weckster BB. Transforming growth factor-beta activity is potentiated by heparin via dissociation of the transforming growth factor beta/alpha 2-macroglobulin inactive complex. *J Cell Biol* 1989;109:441-8.
  26. Ellis SG, Reuben GS, Wilentz J, Douglas JS, King SB. Effect of 18-24 hour heparin administration for prevention of restenosis after uncomplicated coronary angioplasty. *Am Heart J* 1989;117:777-82.
  27. Serruys CW, Baxley WA, Balle TM, Chorro JM, Drans IJ. Early and late effect of heparin infusion following elective angioplasty [abstract]. *Circulation* 1988;78 Suppl II:II-98.
  28. Lehmann KG, Doria RJ, Feuer JM, Hall PK, Hosang DT. Paradoxical increase in restenosis rate with chronic heparin use: final results of a randomised trial [abstract]. *J Am Coll Cardiol* 1991;17 Suppl A:258A.
  29. Hoover RL, Rosenberg RD, Haeering W, Karnovsky M. Inhibition of rat arterial smooth muscle cell proliferation by heparin. II. In vitro studies. *Circ Res* 1980;47:578-83.
  30. Steele PM, Cheshro JH, Stanson AW, et al. Balloon angioplasty: natural history of the pathophysiologic response to injury in a pig model. *Circ Res* 1985;57:105-12.
  31. Gershlick AH, de Bono DP. Restenosis after coronary angioplasty: a proposal of new comparative approaches based on quantitative angiography [letter]. *Br Heart J* 1993;69:469.
  32. de Swiet M, Ward DP, Fidler I. Prolonged heparin therapy in pregnancy causes bone demineralisation. *Br J Obstet Gynaecol* 1983;90:1129-34.
  33. Castellor JJ, Addonizio ML, Rosenberg RD, Karnovsky M. Cultured endothelial cells produce a heparin-like inhibitor of smooth muscle cell growth. *J Cell Biol* 1981;90:372-9.
  34. Serruys FW, Rutsch W, Danchin N, et al. Does the angiotensin converting enzyme inhibitor Cilazapril prevent restenosis after percutaneous transluminal coronary angioplasty? *Circulation* 1992;86:100-10.
  35. Serruys PW, Klein W, Tijssen JGP, et al. Evaluation of ketanserine in the prevention of restenosis after percutaneous transluminal coronary angioplasty: a multicentre randomised double-blind placebo controlled trial. *Circulation* 1993;88:1588-601.

## A RANDOMIZED COMPARISON OF CORONARY-STENT PLACEMENT AND BALLOON ANGIOPLASTY IN THE TREATMENT OF CORONARY ARTERY DISEASE

DAVID L. FISCHMAN, M.D., MARTIN B. LEON, M.D., DONALD S. BAIM, M.D., RICHARD A. SCHATZ, M.D., MICHAEL P. SAVAGE, M.D., IAN PENN, M.D., KATHERINE DETRE, M.D., DR.P.H., LISA VELTRI, M.S., DONALD RICCI, M.D., MASAKIYO NOBUYOSHI, M.D., MICHAEL CLEMAN, M.D., RICHARD HEUSER, M.D., DAVID ALMOND, M.D., PAUL S. TEIRSTEIN, M.D., R. DAVID FISH, M.D., ANTONIO COLOMBO, M.D., JEFFREY BRINKER, M.D., JEFFREY MOSES, M.D., ALEX SHAKNOVICH, M.D., JOHN HIRSHFELD, M.D., STEPHEN BAILEY, M.D., STEPHEN ELLIS, M.D., RANDAL RAKE, B.S., AND SHELDON GOLDBERG, M.D.,  
FOR THE STENT RESTENOSIS STUDY INVESTIGATORS\*

**Abstract Background.** Coronary-stent placement is a new technique in which a balloon-expandable, stainless-steel, slotted tube is implanted at the site of a coronary stenosis. The purpose of this study was to compare the effects of stent placement and standard balloon angioplasty on angiographically detected restenosis and clinical outcomes.

**Methods.** We randomly assigned 410 patients with symptomatic coronary disease to elective placement of a Palmaz-Schatz stent or to standard balloon angioplasty. Coronary angiography was performed at base line, immediately after the procedure, and six months later.

**Results.** The patients who underwent stenting had a higher rate of procedural success than those who underwent standard balloon angioplasty (96.1 percent vs. 89.6 percent,  $P = 0.011$ ), a larger immediate increase in the diameter of the lumen ( $1.72 \pm 0.46$  vs.  $1.23 \pm 0.48$  mm,  $P < 0.001$ ), and a larger luminal diameter immediately after the procedure ( $2.49 \pm 0.43$  vs.  $1.99 \pm 0.47$  mm,  $P < 0.001$ ). At six months, the patients with stented lesions contin-

ued to have a larger luminal diameter ( $1.74 \pm 0.60$  vs.  $1.56 \pm 0.65$  mm,  $P = 0.007$ ) and a lower rate of restenosis (31.6 percent vs. 42.1 percent,  $P = 0.046$ ) than those treated with balloon angioplasty. There were no coronary events (death; myocardial infarction; coronary-artery bypass surgery; vessel closure, including stent thrombosis; or repeated angioplasty) in 80.5 percent of the patients in the stent group and 76.2 percent of those in the angioplasty group ( $P = 0.16$ ). Revascularization of the original target lesion because of recurrent myocardial ischemia was performed less frequently in the stent group than in the angioplasty group (10.2 percent vs. 15.4 percent,  $P = 0.06$ ).

**Conclusions.** In selected patients, placement of an intracoronary stent, as compared with balloon angioplasty, results in an improved rate of procedural success, a lower rate of angiographically detected restenosis, a similar rate of clinical events after six months, and a less frequent need for revascularization of the original coronary lesion. (N Engl J Med 1994;331:496-501.)

THE long-term benefit of coronary balloon angioplasty is limited by the possibility of restenosis of the treated segment, which occurs in approximately 30 to 50 percent of patients.<sup>1-4</sup> Restenosis can be caused by several factors, including elastic recoil of the dilated artery, platelet-mediated thrombus formation, proliferation of smooth-muscle cells, and vascular remodeling.<sup>5</sup> When restenosis develops, it is frequently associated with recurrent myocardial ischemia that necessitates additional revascularization procedures. New approaches to coronary intervention have therefore been developed with the aim of reducing the possibility of restenosis. Debarking coronary atheroma with lasers or atherectomy has not improved the problem of restenosis.<sup>6-9</sup> However, prelimi-

nary evidence suggests that stents may reduce the chance of restenosis by decreasing the elastic recoil of the vessel and sealing intimal flaps, thus providing a wider, smoother coronary lumen.<sup>10,11</sup> To test this hypothesis, we conducted a prospective, randomized trial to compare the rates of restenosis with coronary-stent placement and standard balloon angioplasty.

### METHODS

#### Participating Centers and Investigators

The study centers and investigators were selected on the basis of their experience with implantation of Palmaz-Schatz coronary stents. The study protocol was approved by the institutional review board at each of the 20 centers participating in the trial.

#### Patient Selection

The study population consisted of patients with symptomatic ischemic heart disease and new lesions of the native coronary circulation. The specific angiographic criteria for enrollment included at least 70 percent stenosis, according to the estimate of the investigators; a lesion that was 15 mm or less in length and could be spanned by a single stent; and a vessel diameter of at least 3.0 mm. The criteria for exclusion were a myocardial infarction within the previous seven days; a contraindication to aspirin, dipyridamole, or warfarin sodium; and a left ventricular ejection fraction of 40 percent or less. The angiographic criteria for exclusion were evidence of coronary thrombus, the presence of multiple focal lesions or diffuse disease, serious disease in the left main coronary artery, ostial lesions, and severe vessel tortuosity.

#### Randomization

After the patients had been interviewed to determine their eligibility and had given their informed consent, they were randomly

From Jefferson Medical College, Philadelphia (D.L.F., M.P.S., R.R., S.G.); Washington Cardiology Center, Washington, D.C. (M.B.L.); Beth Israel Hospital, Boston (D.S.B.); Scripps Clinic and Research Center, La Jolla, Calif. (R.A.S., P.S.T.); Victoria General Hospital, Halifax, N.S. (J.P.); the University of Pittsburgh, Pittsburgh (K.D., L.V.); Vancouver General Hospital, Vancouver, B.C. (D.R.); Kokura Memorial Hospital, Kyushu, Japan (M.N.); Yale University, New Haven, Conn. (M.C.); Arizona Heart Institute, Phoenix (R.H.); Toronto General Hospital, Toronto (D.A.); St. Luke's Hospital, Houston (R.D.F.); Centro Cuore Columbus, Milan, Italy (A.C.); Johns Hopkins Hospital, Baltimore (J.B.); Lenox Hill Hospital, New York (J.M., A.S.); Hospital of the University of Pennsylvania, Philadelphia (J.H.); the University of Texas at San Antonio, San Antonio (S.B.); and the Cleveland Clinic Foundation, Cleveland (S.E.). Address reprint requests to Dr. Goldberg at Jefferson Medical College, Division of Cardiology, Suite 403, 1025 Walnut St., Philadelphia, PA 19107.

Supported in part by a grant from Johnson and Johnson Interventional Systems.

\*Additional participants in the Stent Restenosis Study (STRESS) trial are listed in the Appendix.



assigned to either stent placement or balloon angioplasty. Randomization of the patients, stratified according to center with a block design, was carried out by means of sealed envelopes. The randomization sequence was developed so that an equal number of patients would be assigned to each treatment at each center.

### Procedural Protocol

#### Stent Placement

The Palmaz-Schatz stent is composed of two rigid 7-mm slotted stainless-steel tubes connected by a 1-mm central bridging strut (Johnson and Johnson Interventional Systems, Warren, N.J.). The stent, which is 1.6 mm in diameter in the unexpanded state, is mounted on a balloon catheter and protected by an outer sheath during passage to the target site. When the sheath is withdrawn, inflation of the balloon catheter expands the stent. Technical details of the design and placement of the Palmaz-Schatz coronary stent have been described elsewhere.<sup>12,13</sup>

Patients assigned to stent placement received nonenteric aspirin (325 mg daily), dipyridamole (75 mg three times a day), and treatment with a calcium-channel antagonist, initiated at least 24 hours before the procedure. In addition, patients received intravenous low-molecular-weight dextran (dextran 40, given at a dose of 100 ml per hour for two hours before stenting and at a dose of 50 ml per hour during and after the procedure, for a total volume of 1 liter). During the procedure, patients received an initial bolus injection of heparin (10,000 to 15,000 units) supplemented as needed to maintain an activated clotting time of more than 300 seconds. The heparin infusion was discontinued at the termination of the procedure and reinstituted four to six hours after hemostasis of the site of vascular access had been achieved. Warfarin sodium was begun on the day of the procedure. Heparin and warfarin sodium were both administered for at least 72 hours or until a prothrombin time of 16 to 18 seconds had been achieved (international normalized ratio, 2.0 to 3.5). After patients were discharged from the hospital, dipyridamole and warfarin sodium were continued for one month, and aspirin was continued indefinitely.

#### Angioplasty Protocol

Angioplasty was performed with the use of conventional techniques. Aspirin was prescribed, but warfarin sodium was not administered. Investigators attempted to achieve an optimal result with balloon angioplasty, which was defined as residual stenosis of less than 30 percent of the luminal diameter, according to a visual estimate. A crossover to stent placement was permitted as a "bail-out" procedure in the case of abrupt or threatened closure, defined as a dissection of the artery with compromised antegrade blood flow (Thrombolysis in Myocardial Infarction [TIMI] grade, <3) or persistent stenosis of over 50 percent of the luminal diameter in association with evidence of myocardial ischemia (chest pain, electrocardiographic changes, or both).

#### Follow-up

Patients were required to have clinical follow-up studies after one, three, and six months. Coronary angiography was required at six months in all the patients except those who had died or undergone coronary-artery bypass surgery or repeated angioplasty for abrupt closure during the first 14 days after the initial revascularization. Angiography performed before four months was allowed on the basis of clinical indications. However, if restenosis was not found, a subsequent angiogram was obtained after four months.

#### Angiographic Analysis

Angiography was performed in two orthogonal views. Intracoronary nitroglycerin (200 mg) was injected before all angiographic assessments. Angiograms were analyzed at the Core Angiographic Laboratory at Jefferson Medical College. Quantitative analysis was performed with the use of a validated edge-detection algorithm.<sup>14</sup> Vessel edges were determined with the computerized algorithm, and luminal diameters were measured with the dye-filled catheter as a reference. The diameters of the normal segments proximal and

distal to the treated area were averaged to determine the reference diameter. The minimal luminal diameter, reference diameter, and percentage of stenosis were calculated as the mean values from two orthogonal projections. The percentage of elastic recoil was defined as the largest inflated-balloon diameter minus the postprocedural minimal luminal diameter divided by the inflated-balloon diameter. In addition, coronary lesions were assessed for eccentricity, calcification, thrombus, plaque ulceration, tortuosity, and postprocedural dissection. Definitions used for this morphologic analysis and prior validation studies of the quantitative angiographic analysis have been described elsewhere.<sup>11,13,15</sup>

#### End Points

The primary end point of the trial was angiographic evidence of restenosis, defined as at least 50 percent stenosis on the follow-up angiogram. Secondary angiographic end points included angiographic evidence of procedural success and the absolute minimal luminal diameter after the procedure and at follow-up. Angiographic evidence of procedural success was defined as a reduction in stenosis to 50 percent or less by quantitative analysis.

Clinical evidence of procedural success was defined as angiographic evidence of success without a major complication (death, myocardial infarction, or coronary-artery bypass surgery) during the index hospitalization. The secondary clinical end point was a composite end point, defined as whichever of the following occurred first: death, myocardial infarction, coronary bypass surgery, or the need for repeated angioplasty within the first 6 months ( $\pm 60$  days) after the initial revascularization. Myocardial infarction was documented by the presence of new Q waves of at least 0.04 second's duration or a creatine kinase level or MB fraction at least twice the upper limit of normal. Clinical events were classified as early (occurring from day 0 to day 14) or late (occurring after 14 days). Revascularization of the target lesion was defined as angioplasty or bypass surgery performed because of restenosis of the target lesion in association with recurrent angina, objective evidence of myocardial ischemia, or both. Other events included abrupt vessel closure (after the patient had left the catheterization laboratory) and hemorrhagic complications, defined as a cerebrovascular accident, bleeding requiring transfusion, or the need for vascular surgery.

Clinical and angiographic data were forwarded to the Data Coordinating Center at the University of Pittsburgh for statistical analyses. Adverse events were audited and reviewed by members of the Steering Committee. The primary analysis of angiographic and procedural outcomes was based on the intention-to-treat principle. We also performed a secondary analysis of the rate of restenosis according to the treatment received.

For the analysis of continuous data, two-tailed t-tests were used to assess differences between the two treatment groups. The results are expressed as means  $\pm$  SD. Categorical data, which are presented as rates, were compared by chi-square test, except for the composite clinical end point and revascularization of the target lesion, which were analyzed by means of Kaplan-Meier survival curves, with differences between the two treatment groups compared by Wilcoxon test.<sup>16</sup> Multiple linear regression was used to assess the relation between the luminal diameter at follow-up and multiple clinical and angiographic variables, including age, sex, location of the lesion, vessel diameter, and postprocedural luminal diameter.

### RESULTS

Between January 1991 and February 1993, 410 patients were enrolled in the study; 207 patients were randomly assigned to stent placement, and 203 to angioplasty. After randomization, three patients (two in the stent group and one in the angioplasty group) were excluded because they did not meet all the enrollment criteria. Thus, the final study group comprised 407 patients. Their base-line clinical and angiographic characteristics are shown in Table 1. More

men were assigned to the stent group than to the angioplasty group, and the patients in the stent group had lesions that were slightly longer, with a higher incidence of eccentricity, but the two groups were well matched with respect to other clinical characteristics.

#### Procedural and Early Clinical Outcome

Stents were placed in 197 of the 205 patients (96.1 percent) randomly assigned to this therapy. One patient, in whom stent placement failed because of an inability to cross the lesion with a guide wire, was treated medically. Seven patients were switched to angioplasty: three because of an inability to place the stent and four because of lesion characteristics deemed unfavorable for stent placement at the time of the procedure. In the angioplasty group, six patients required emergency coronary-artery bypass surgery. In addition, 15 patients were switched to alternative therapies: 14 (6.9 percent) to emergency stent place-

Table 1. Base-Line Clinical and Angiographic Characteristics of Patients Assigned to Stent Placement or Angioplasty.\*

CHARACTERISTIC	STENT GROUP (N = 205)	ANGIOPLASTY GROUP (N = 202)
Male — % of patients	83	73†
Age — yr	60±10	60±10
Diabetes — % of patients	15	16
Hypertension — % of patients	43	45
Hyperlipidemia — % of patients	44	48
Current smoker — % of patients	21	24
History of myocardial infarction — % of patients	37	36
Recent myocardial infarction (within previous 6 wk) — % of patients	18	15
Unstable angina — % of patients	47	48
Pain at rest	33	39
Pain with electrocardiographic changes	23	26
Postinfarction angina	7	6
No. of diseased vessels — % of patients		
1	64	68
2	27	21
3	9	11
Ejection fraction — %	61±12	61±11
Target vessel — % of patients		
Left anterior descending	47	48
Left circumflex	16	13
Right coronary artery	37	39
Calcification — % of patients	17	15
Thrombus — % of patients		
Definite	2	1
Possible	15	9
Eccentricity — % of patients	66	54‡
Lesion angulation >45° — % of patients	13	18
Lesion length — mm	9.6±3.0	8.7±2.7§
Stenosis — % of luminal diameter	75±9	75±8

\*Plus-minus values are means ±SD.

†P<0.05.

‡P = 0.02.

§P<0.001.

Table 2. Procedural Outcomes and Clinical Events.

VARIABLE	STENT GROUP (N = 205)	ANGIOPLASTY GROUP (N = 202)	P VALUE
% of patients			
Procedural outcome			
Angiographic success	99.5	96.5	0.04
Reading at study center	99.5	92.6	<0.001
Quantitative analysis	96.1	89.6	0.011
Clinical success			
Early events (0–14 days)			
Death	0	1.5	0.12
Myocardial infarction/Q wave	5.4/2.9	5.0/3.0	0.85/1.0
Coronary bypass surgery	2.4	4.0	0.38
Abrupt closure*	3.4	1.5	0.34
Repeated angioplasty	2.0	1.0	0.69
Any event	5.9	7.9	0.41
Late events (15–240 days)			
Death	1.5	0	0.25
Myocardial infarction/Q wave	1.5/1.0	2.0/0.5	0.72/1.0
Coronary bypass surgery	2.4	4.5	0.26
Repeated angioplasty	9.8	11.4	0.59
Target-vessel revascularization	10.2	15.4	0.06
Any event	15.1	15.8	0.84
All events (0–240 days)			
Death	1.5	1.5	0.99
Myocardial infarction/Q wave	6.3/3.4	6.9/3.5	0.81/0.5
Coronary bypass surgery	4.9	8.4	0.15
Repeated angioplasty	11.2	12.4	0.72
Any event	19.5	23.8	0.16
Bleeding and vascular complications			
Cerebrovascular accident	1.0	0.5	1.0
Surgical vascular repair	3.9	2.0	0.25
Bleeding requiring transfusion	4.9	2.5	0.11
Any event	7.3	4.0	0.14

\*After the patient left the catheterization laboratory.

ment as a bailout procedure (1 of the 14 subsequent required emergency bypass surgery) and 1 to directional atherectomy.

Procedural and early clinical outcomes are shown in Table 2. According to the quantitative analysis there was angiographic evidence of procedural success in 204 of the 205 patients (99.5 percent) randomly assigned to undergo stent placement and in 187 of the 202 patients (92.6 percent) randomly assigned to undergo angioplasty (P<0.001). The clinical success rates were 96.1 percent and 89.6 percent, respectively (P = 0.011).

Abrupt vessel closure occurred in 10 patients after they had left the catheterization laboratory: 7 in the stent group and 3 in the angioplasty group (3.4 and 1.5 percent, respectively; P = 0.34). In the three patients in the angioplasty group, the closure occurred after the stent had been placed as a bailout measure. Abrupt closure occurred an average of 6 days (range 2 to 14) after the procedure, and in 6 of the patients, it occurred after hospital discharge. The patients with abrupt closures had major cardiac events (two died and eight had nonfatal myocardial infarctions). The proportions of patients with any major cardiac event (death, myocardial infarction, coronary bypass surgery, or repeated angioplasty within 14 days after the procedure) were 5.9 percent in the stent group and 7.9 percent in the angioplasty group.

(Table 2). Bleeding and vascular complications occurred more commonly in the stent group than in the angioplasty group (7.3 percent vs. 4.0 percent,  $P = 0.14$ ). The hospital stay after the procedure was longer in the stent group (5.8 days vs. 2.8 days,  $P < 0.001$ ).

#### Angiographic Results

Angiography was repeated at six months in 336 of the 383 patients (88 percent) eligible for follow-up. Angiography was not repeated in 28 patients in the stent group because of refusal (15 patients) or ineligibility due to stent thrombosis (7), death (3), early coronary bypass surgery (2), or inability to perform the study procedures (1). In the angioplasty group, 43 patients did not have follow-up angiography because of refusal (32) or ineligibility due to early coronary bypass surgery (7), abrupt vessel closure (3), or death (1). The rate of restenosis was 31.6 percent (56 of 177 patients) in the stent group and 42.1 percent (67 of 159) in the angioplasty group ( $P = 0.046$ ). The rates of restenosis among the patients who received their assigned therapy were 30.0 percent in the stent group and 43.0 percent in the angioplasty group ( $P = 0.016$ ).

The luminal dimensions at base line, immediately after the procedure, and at follow-up are shown in Table 3. At base line, there was no difference in the reference diameter or the severity of stenosis between the two groups. After the procedure, a larger immediate gain in the luminal diameter was achieved in the patients who underwent stent placement than in those who underwent angioplasty, resulting in a larger mean ( $\pm$ SD) diameter in the stent group ( $2.49 \pm 0.43$  vs.  $1.99 \pm 0.47$  mm,  $P < 0.001$ ). At follow-up, the stent group had a larger mean reduction in the luminal diameter ( $0.74 \pm 0.58$  vs.  $0.38 \pm 0.66$  mm,  $P < 0.001$ ) but a larger net gain, resulting in a larger luminal diameter at follow-up ( $1.74 \pm 0.60$  vs.  $1.56 \pm 0.65$  mm,  $P = 0.007$ ). These data are shown in Figure 1. A step-wise linear regression analysis showed that the luminal diameter immediately after the procedure was the most important predictor of the luminal diameter at six months ( $b = 0.41$ ,  $P < 0.001$ ), irrespective of the procedure used. Additional important determinants included a larger reference diameter ( $b = 0.31$ ,  $P < 0.001$ ) and location of the lesion in a vessel other than the left anterior descending coronary artery ( $b = 0.14$ ,  $P = 0.029$ ).

#### Late Clinical Follow-up

Data on late cardiac events and all events are shown in Table 2. Clinical follow-up data were available for 406 of the 407 patients. Although the numbers of patients who died or had myocardial infarctions were comparable in the two groups, fewer patients in the stent group underwent revascularization of the target lesion (10.2 percent vs. 15.4 percent,  $P = 0.06$ ) (Fig. 2). Event-free survival was 80.5 percent in the stent

Table 3. Angiographic Results in the Stent and Angioplasty Groups.\*

VARIABLE	STENT GROUP (N = 205)	ANGIOPLASTY GROUP (N = 202)	P VALUE
Before the procedure			
Reference vessel (mm)	$3.03 \pm 0.42$	$2.99 \pm 0.50$	0.30
Minimal luminal diameter (mm)	$0.77 \pm 0.27$	$0.75 \pm 0.25$	0.48
Stenosis (% of luminal diameter)	$75 \pm 9$	$75 \pm 8$	0.81
After the procedure			
Reference vessel (mm)	$3.05 \pm 0.40$	$2.99 \pm 0.46$	0.15
Minimal luminal diameter (mm)	$2.49 \pm 0.43$	$1.99 \pm 0.47$	$< 0.001$
Stenosis (% of luminal diameter)	$19 \pm 11$	$35 \pm 14$	$< 0.001$
Elastic recoil (%)	$15 \pm 11$	$24 \pm 15$	$< 0.001$
Dissection (% of patients)	7	25	$< 0.001$
At follow-up			
Reference vessel (mm)	$3.00 \pm 0.41$	$2.98 \pm 0.49$	0.74
Minimal luminal diameter (mm)	$1.74 \pm 0.60$	$1.56 \pm 0.65$	0.007
Stenosis (% of luminal diameter)	$42 \pm 18$	$49 \pm 19$	0.001
Restenosis (% of patients)	31.6	42.1	0.046
Change in minimal luminal diameter			
Immediate gain (mm)	$1.72 \pm 0.46$	$1.23 \pm 0.48$	$< 0.001$
Late loss (mm)	$0.74 \pm 0.58$	$0.38 \pm 0.66$	$< 0.001$
Net gain (mm)	$0.98 \pm 0.62$	$0.80 \pm 0.63$	0.01

\*Plus-minus values are means  $\pm$ SD. Immediate gain refers to the minimal luminal diameter immediately after the procedure minus the diameter before the procedure. Late loss refers to the minimal luminal diameter immediately after the procedure minus the diameter at follow-up. Net gain refers to the minimal luminal diameter at follow-up minus the diameter before the procedure.

group, as compared with 76.2 percent in the angioplasty group ( $P = 0.16$ ) (Fig. 3). Among the patients eligible for follow-up, a larger proportion of those in the stent group remained free of angina (78.9 percent vs. 71.1 percent,  $P = 0.076$ ).

#### DISCUSSION

In this trial, we compared stent placement with balloon angioplasty for the treatment of new focal coronary stenoses in larger vessels; we found a reduc-




Figure 1. Minimal Diameter of the Lumen at Base Line, Immediately after Stent Placement or Angioplasty, and at Follow-up.

There was no difference in base-line values between the stent and angioplasty groups. Immediately after the procedure, the patients in the stent group had a larger minimal luminal diameter than those in the angioplasty group. Six months later, both groups had reduced values, and a significant difference in diameter persisted between the two groups.

tion in the rate of angiographic restenosis at six months with the stenting procedure. This reduction was associated with a reduction in the need for repeat revascularization due to ischemia-associated restenosis.

Our findings contrast with those of previous investigations that examined the efficacy of pharmacologic agents in preventing restenosis.<sup>17-24</sup> Of the newer interventional procedures, only directional atherectomy has been subjected to careful prospective, randomized studies to assess its efficacy in reducing restenosis, as compared with the efficacy of angioplasty.<sup>7,8</sup> Those studies showed either no benefit of atherectomy or a minimal reduction in restenosis with more frequent major complications.

Like the Coronary Angioplasty versus Excisional Atherectomy Trial (CAVEAT),<sup>7</sup> our study shows that the most important determinant of the luminal diameter at six months was the luminal diameter achieved immediately after the procedure. It seems plausible that the reduction in restenosis in our stent group was due to the significantly larger luminal diameter obtained immediately after placement of the stent, as compared with the luminal diameter immediately after angioplasty. The residual stenosis in the stent group (19 percent) was roughly half that in the angioplasty group (35 percent) and 10 percentage points less than the residual stenosis in patients undergoing directional atherectomy.<sup>7</sup> Although the larger immediate gain in luminal diameter was offset by a larger subsequent loss, the net gain remained larger in the patients in the stent group (Fig. 1). Multivariate analysis showed that the luminal diameter immediately after the procedure was the most powerful predictor of the luminal diameter at follow-up, regardless of whether stenting or balloon angioplasty achieved this result. Therefore, it was not the specific technique used, but rather its efficacy in achieving a larger lu-




Figure 2. Kaplan-Meier Curves for Revascularization of the Target Lesion.

Fewer patients in the stent group than in the angioplasty group required revascularization of the target lesion because of ischemia.




Figure 3. Kaplan-Meier Survival Curves for Major Cardiac Events (Death, Myocardial Infarction, Coronary-Artery Bypass Surgery, and Repeated Angioplasty).

minal diameter that was the determining factor, an idea that has been suggested previously.<sup>25</sup> In addition, stenting resulted in a larger diameter with less risk of intimal disruption and elastic recoil, thereby acting as an effective intravascular scaffold. The ability of the stent to serve as a scaffold was further demonstrated in the 14 patients in the angioplasty group (6.9 percent) who were switched to stent placement for treatment of imminent or actual closure after balloon angioplasty had failed. At the inception of this trial, stent placement as a bailout measure, which at the time was not available as a routine procedure, was considered equivalent to emergency coronary-artery bypass surgery. Thirteen of the 14 patients who underwent stent placement as a bailout measure had balloon-induced dissections or luminal compromise associated with chest pain or electrocardiographic changes, suggesting that these patients would have had serious clinical events if stent placement had not been available. Therefore, the availability of stent placement probably decreased the rate of clinical events in the angioplasty group. This study thus represents a comparison of two treatment strategies: elective stent placement and elective balloon angioplasty with stent placement available as a bailout measure.

Several limitations of stent placement need to be emphasized. Stent thrombosis occurred in 3.4 percent of the patients who underwent stent placement as an elective procedure and in 21.4 percent of those in whom stent placement was used as a bailout technique. These thrombotic events occurred 2 to 14 days after placement of the stent, with six instances of thrombosis after discharge, and invariably resulted in major clinical complications. Furthermore, the intense anticoagulation and antiplatelet regimen associated with stent placement resulted in nearly twice the number of hemorrhagic and peripheral vascular complications associated with angioplasty, as well as a prolonged hospital stay.

Although the frequency with which follow-up angiography was performed was relatively high in both

groups, there was a higher rate of angiographic follow-up in the stent group (92 percent vs. 83 percent,  $P = 0.008$ ). This difference, which may bias the rate of restenosis in favor of stent placement, is a limitation of the study.

In conclusion, elective stent placement, as compared with angioplasty, has a higher clinical success rate and reduces the incidence of restenosis and the need for subsequent revascularization of the treated lesion. The reduction in restenosis is not associated with an increase in major cardiac events, despite the limitations imposed by stent thrombosis and hemorrhagic complications. The use of antithrombotic stent coatings, improved techniques to optimize expansion of the stent during implantation, and compression and closure devices at the site of arteriotomy may address these limitations. If they are effectively overcome, implantation of the Palmaz-Schatz stent may become the preferred treatment in selected patients with new lesions in large coronary arteries.

#### APPENDIX

The following institutions and investigators participated in the STRESS trial: Arizona Heart Institute, Phoenix (E. Davis, W. Catran, and K. Waters); Beth Israel Hospital, Boston (D.J. Diver, J. Carrozza, and C. Senerchia); Centro Cuore Columbus, Milan, Italy (Y. Almagor and M. Bernati); Cleveland Clinic Foundation, Cleveland (P. Whitlow); Florida Heart Hospital, Orlando (C. Curry, C.B. Saenz, W.H. Willis, Jr., R.J. Ivanhoe, and N. Granger); Hospital of the University of Pennsylvania, Philadelphia (H. Herman, D. Kolansky, W. Laskey, and D. DiAngelo); Johns Hopkins Hospital, Baltimore (V. Coombs); Lenox Hill Hospital, New York (E.M. Kreps, J. Strain, N. Cohen, J. Higgins, and C. Udemir); Scripps Clinic and Research Foundation, San Diego, Calif. (N. Morris and M. Dowling); St. Luke's Hospital, Houston (M. Harlan and B. Lambert); Thomas Jefferson University Hospital, Philadelphia (A. Zalewski, P. Walinsky, and D. Porter); Toronto General Hospital, Toronto (L. Lazzam, C. Lazzam, and P. Slaughter); University of Texas at San Antonio, San Antonio (J.P. Hennecken, S. Kiesz, and A. Briscoe); Vancouver General Hospital, Vancouver, B.C. (C.E. Buller and A. McCarthy); Victoria General Hospital, Halifax, N.S. (B. O'Neil, C.J. Foster, C.M. Peck, K.A. Foshay, and N.L. Fitzgerald); Victoria Hospital, London, Ont. (N. Murray-Parson and L. Marziali); Washington Cardiology Center, Washington, D.C. (K. Donovan); Yale University, New Haven, Conn. (H.S. Cabin and R.E. Rosen); *Data Coordinating Center*: Department of Epidemiology, University of Pittsburgh, Pittsburgh (K. Detre, V. Niedermyer, L. Kennard, and L. Vetri); *Core Angiographic Laboratory*: Thomas Jefferson University Hospital, Philadelphia (R. Rake, S. Gebhardt, D.L. Fischman, M.P. Savage, and S. Goldberg); *Steering Committee*: D.S. Baim, S. Goldberg, M.B. Leon, I. Penn, and R.A. Schatz.

#### REFERENCES

1. Holmes DR Jr, Vlietstra RE, Smith HC, et al. Restenosis after percutaneous transluminal coronary angioplasty (PTCA): a report from the PTCA Registry of the National Heart, Lung, and Blood Institute. *Am J Cardiol* 1984;53:77C-81C.
2. Gruentzig AR, King SB III, Schlumpf M, Siegenthaler W. Long-term follow-up after percutaneous transluminal coronary angioplasty: the early Zurich experience. *N Engl J Med* 1987;316:1127-32.
3. Nobuyoshi M, Kimura T, Nosaka H, et al. Restenosis after successful percutaneous transluminal coronary angioplasty: serial angiographic follow-up of 229 patients. *J Am Coll Cardiol* 1988;12:616-23.
4. Hirshfeld JW Jr, Schwartz JS, Jugo R, et al. Restenosis after coronary angioplasty: a multivariate statistical model to relate lesion and procedure variables to restenosis. *J Am Coll Cardiol* 1991;18:647-56.
5. Waller BF. "Crackers, breakers, stretchers, drillers, scrapers, shavers, burners, welders and melters" — the future treatment of atherosclerotic coronary artery disease? A clinical-morphologic assessment. *J Am Coll Cardiol* 1989;13:969-87.
6. Bittl JA, Sanborn TA, Tchong JE, Siegel RM, Ellis SG. Clinical success, complications and restenosis rates with excimer laser coronary angioplasty. *Am J Cardiol* 1992;70:1533-9.
7. Topol EJ, Leya F, Pinkerton CA, et al. A comparison of directional atherectomy with coronary angioplasty in patients with coronary artery disease. *N Engl J Med* 1993;329:221-7.
8. Adelman AG, Cohen EA, Kimball BP, et al. A comparison of directional atherectomy with balloon angioplasty for lesions of the left anterior descending coronary artery. *N Engl J Med* 1993;329:228-33.
9. Safian RD, Niaz KA, Strzelecki M, et al. Detailed angiographic analysis of high-speed mechanical rotational atherectomy in human coronary arteries. *Circulation* 1993;88:961-8.
10. Ellis SG, Savage M, Fischman D, et al. Restenosis after placement of Palmaz-Schatz stents in native coronary arteries: initial results of a multicenter experience. *Circulation* 1992;86:1836-44.
11. Savage M, Fischman D, Leon M, et al. Long-term angiographic and clinical outcome after implantation of balloon-expandable stents in the native coronary circulation. *J Am Coll Cardiol* (in press).
12. Schatz RA, Baim DS, Leon M, et al. Clinical experience with the Palmaz-Schatz coronary stent: initial results of a multicenter study. *Circulation* 1991;83:148-61.
13. Fischman DL, Savage MP, Ellis S, et al. The Palmaz-Schatz stent. In: Reiber JHC, Serruys PW, eds. *Advances in quantitative coronary arteriography*. Vol. 137 of *Developments in cardiovascular medicine*. Dordrecht, the Netherlands: Kluwer Academic, 1993:553-66.
14. Mancini GBJ, Simon SB, McGillem MJ, LeFree MT, Friedman HZ, Vogel RA. Automated quantitative coronary arteriography: morphologic and physiologic validation in vivo of a rapid digital angiographic method. *Circulation* 1987;75:452-60. [Erratum, *Circulation* 1987;75:1199.]
15. Fischman DL, Savage MP, Leon MB, et al. Effect of intracoronary stenting on intimal dissection after balloon angioplasty: results of quantitative and qualitative coronary analysis. *J Am Coll Cardiol* 1991;18:1445-51.
16. Kalbfleisch JD, Prentice RL. *The statistical analysis of failure time data*. New York: John Wiley, 1980.
17. Pepine CJ, Hirshfeld JW, Macdonald RG, et al. A controlled trial of corticosteroids to prevent restenosis after coronary angioplasty. *Circulation* 1990;81:1753-61.
18. Thornton MA, Gruentzig AR, Hollman J, King SB III, Douglas JS. Coumadin and aspirin in prevention of recurrence after transluminal coronary angioplasty: a randomized study. *Circulation* 1984;69:721-7.
19. Schwartz L, Bourassa MG, Lesperance J, et al. Aspirin and dipyridamol in the prevention of restenosis after percutaneous transluminal coronary angioplasty. *N Engl J Med* 1988;318:1714-9.
20. Ellis SG, Roubin GS, Wilentz J, Douglas JS Jr, King SB III. Effect of 18- to 24-hour heparin administration for prevention of restenosis after uncomplicated coronary angioplasty. *Am Heart J* 1989;117:777-82.
21. Dehmer GJ, Popma JJ, van den Berg EK, et al. Reduction in the rate of early restenosis after coronary angioplasty by a diet supplemented with n-3 fatty acids. *N Engl J Med* 1988;319:733-40.
22. Reis GJ, Boucher TM, Sipperly ME, et al. Randomized trial of fish oil for prevention of restenosis after coronary angioplasty. *Lancet* 1989;2:177-81.
23. Whitworth HB, Roubin GS, Hollman J, et al. Effect of nifedipine on recurrent stenosis after percutaneous transluminal coronary angioplasty. *J Am Coll Cardiol* 1986;8:1271-6.
24. Corcos T, David PR, Val PG, et al. Failure of diltiazem to prevent restenosis after percutaneous transluminal coronary angioplasty. *Am Heart J* 1985;109:926-31.
25. Kuntz RE, Gibson CM, Nobuyoshi M, Baim DS. Generalized model of restenosis after conventional balloon angioplasty, stenting and directional atherectomy. *J Am Coll Cardiol* 1993;21:15-25.

# TEXTBOOK OF INTERVENTIONAL CARDIOLOGY

5TH EDITION

**Eric J. Topol, MD**

*Director, Scripps Translational Science Institute*

*Chief Academic Officer, Scripps Health*


*Professor of Translational Genomics, The Scripps Research Institute*

*Senior Consultant, Division of Cardiovascular Diseases*

*Scripps Clinic*

*La Jolla, California*

**SAUNDERS**



**SAUNDERS**  
ELSEVIER

1600 John F. Kennedy Blvd.  
Ste 1800  
Philadelphia, PA 19103-2899

TEXTBOOK OF INTERVENTIONAL CARDIOLOGY

ISBN: 978-1-4160-4835-0

Copyright © 2008, 2003, 1999, 1994, 1990 by Saunders, an imprint of Elsevier Inc.

**All rights reserved.** No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher. Permissions may be sought directly from Elsevier's Rights Department: phone: (+1) 215 239 3804 or (+44) 1865 843830 (UK); fax: (+44) 1865 853333; e-mail: [healthpermissions@elsevier.com](mailto:healthpermissions@elsevier.com). You may also complete your request on-line via the Elsevier website at <http://www.elsevier.com/permissions>.

#### Notice

Knowledge and best practice in this field are constantly changing. As new research and experience broaden our knowledge, changes in practice, treatment, and drug therapy may become necessary or appropriate. Readers are advised to check the most current information provided (i) on procedures featured or (ii) by the manufacturer of each product to be administered, to verify the recommended dose or formula, the method and duration of administration, and contraindications. It is the responsibility of the practitioner, relying on his or her experience and knowledge of the patient, to make diagnoses, to determine dosages and the best treatment for each individual patient, and to take all appropriate safety precautions. To the fullest extent of the law, neither the Publisher nor the Editor assumes any liability for any injury and/or damage to persons or property arising out of or related to any use of the material contained in this book.

The Publisher

#### Library of Congress Cataloging-in-Publication Data

Textbook of interventional cardiology / edited by Eric J. Topol.—5th ed.

p. ; cm

Includes bibliographical references and index.

ISBN 978-1-4160-4835-0

1. Heart—Interventional radiology. 2. Angioplasty. 3. Cardiovascular system—Diseases—Treatment. I. Topol, Eric J., 1954-

[DNLM: 1. Cardiovascular Diseases—surgery. 2. Cardiovascular Diseases—drug therapy. 3. Cardiovascular Surgical Procedures—methods. WG 168 T355.2008]

RD598.35.J55T49 2008

617.4'12059—dc22

2007018033

*Executive Publisher:* Natasha Andjelkovic  
*Developmental Editor:* Agnes Hunt Byrne  
*Publishing Services Manager:* Frank Polizzano  
*Project Manager:* Michael H. Goldberg  
*Design Direction:* Steven Stave

Printed in China

Last digit is the print number: 9 8 7 6 5 4 3 2

Working together to grow  
libraries in developing countries

[www.elsevier.com](http://www.elsevier.com) | [www.bookaid.org](http://www.bookaid.org) | [www.sabre.org](http://www.sabre.org)

ELSEVIER

BOOKAID  
International

Sabre Foundation

A1274